Lecture 1

Asymptotic behaviour of random Lipschitz functions

Last updated: 2026/03/31

Lecturer: Ron Peled

Scribe: Ofir Schnabel

Tel Aviv University

Spring 2026

These notes are based on the lectures given by Ron Peled in Tel Aviv University during the spring semester of 2026. They were written by Ofir Schnabel. Please send comments and corrections to ofirschnabel@mail.tau.ac.il.

Introduction

We start with a brief introduction to the course.

Definition 1.1 (Lipschitz functions)

Let $(X,d_X)$ and $(Y,d_Y)$ be metric spaces. A function $f: X \to Y$ is called $L$-Lipschitz if $d_Y(f(x),f(y)) \le L\, d_X(x,y)$ for all $x,y \in X$.

Remark 1.2

(1) Any $L$-Lipschitz function is uniformly continuous.

(2) Rademacher's theorem: If $X = \mathbb{R}^n$ and $Y = \mathbb{R}^m$ then any $L$-Lipschitz function $f$ is differentiable almost everywhere (with respect to Lebesgue measure) and $\|\nabla f\| \le L$ wherever $\nabla f$ exists.

(3) Kirszbraun's theorem: If $X$ is any subset of a Hilbert space $H$ and $Y = H$ then any $L$-Lipschitz function $f: X \to Y$ can be extended to an $L$-Lipschitz function $\tilde{f}: H \to H$.

(4) The Lipschitz constant of a function $f: X \to Y$ is the smallest $L$ such that $f$ is $L$-Lipschitz. It is denoted $\mathrm{Lip}(f)$.

Exercise 1.3

Let $f: X \to Y$ and $g: Y \to Z$ be Lipschitz functions between metric spaces. Show that $\mathrm{Lip}(g \circ f) \le \mathrm{Lip}(g)\,\mathrm{Lip}(f)$.

Solution

Let $L = \mathrm{Lip}(f)$ and $M = \mathrm{Lip}(g)$. For any $x,y \in X$ we have $d_Y(f(x),f(y)) \le L\, d_X(x,y)$ and $d_Z(g(f(x)),g(f(y))) \le M\, d_Y(f(x),f(y))$. Combining these gives $d_Z(g(f(x)),g(f(y))) \le ML\, d_X(x,y)$, so $g \circ f$ is $ML$-Lipschitz.

Exercise 1.4

Let $f: \mathbb{R}^n \to \mathbb{R}$ be a differentiable function. Show that $\mathrm{Lip}(f) = \sup_{x \in \mathbb{R}^n} \|\nabla f(x)\|$.

Solution

Suppose $f$ is $L$-Lipschitz. Then for any $x \in \mathbb{R}^n$ and $v \in \mathbb{R}^n$ with $\|v\| = 1$ we have $|f(x+tv) - f(x)| \le L|t|$, so $|\frac{f(x+tv)-f(x)}{t}| \le L$. Taking $t \to 0$ gives $|\nabla_v f(x)| \le L$, where $\nabla_v f(x) = \nabla f(x) \cdot v$ is the directional derivative. Choosing $v = \nabla f(x)/\|\nabla f(x)\|$ (when $\nabla f(x) \neq 0$) yields $\|\nabla f(x)\| \le L$. Conversely, if $\|\nabla f(x)\| \le L$ for all $x$, then by the mean value theorem $|f(x) - f(y)| \le L\|x-y\|$, so $f$ is $L$-Lipschitz.

Exercise 1.5

Let $f: \mathbb{R}^n \to \mathbb{R}$ be a convex function. Show that $f$ is Lipschitz if and only if it is bounded on some open set.

Solution

If $f$ is $L$-Lipschitz then it is bounded on any bounded set. Conversely, suppose $f \le M$ on the open ball $B(x_0,r)$. For any $x \neq y$, let $z = y + \frac{r}{2}\frac{y-x}{\|y-x\|}$. Then $z \in B(x_0,r)$ and $y = \frac{2}{r}\|y-x\| z + (1 - \frac{2}{r}\|y-x\|) x$. By convexity, $f(y) \le \frac{2}{r}\|y-x\| f(z) + (1 - \frac{2}{r}\|y-x\|) f(x)$, so $f(y) - f(x) \le \frac{2}{r}\|y-x\|(f(z) - f(x)) \le \frac{4M}{r}\|y-x\|$. Swapping $x$ and $y$ gives $|f(x) - f(y)| \le \frac{4M}{r}\|x-y\|$, so $f$ is $\frac{4M}{r}$-Lipschitz.

Exercise 1.6

Let $f: \mathbb{R}^n \to \mathbb{R}$ be a convex function. Show that $\mathrm{Lip}(f) = \max_{x \in \mathbb{R}^n} \max_{g \in \partial f(x)} \|g\|$, where $\partial f(x) = \{g : f(y) \ge f(x) + g \cdot (y-x) \text{ for all } y\}$ is the subdifferential of $f$ at $x$.

Solution

Suppose $f$ is $L$-Lipschitz. Then for any $x$ and $g \in \partial f(x)$, taking $y = x + \frac{g}{\|g\|}$ gives $\|g\| = g \cdot (y-x) \le f(y) - f(x) \le L\|y-x\| = L$. Conversely, if $\|g\| \le L$ for all $g \in \partial f(x)$ and all $x$, then $f(y) \ge f(x) + g \cdot (y-x) \ge f(x) - L\|y-x\|$, and swapping $x$ and $y$ gives $|f(x) - f(y)| \le L\|x-y\|$.

Exercise 1.7

Let $f: \mathbb{R}^n \to \mathbb{R}$ be a convex function. Show that $f$ is Lipschitz if and only if $\sup_{x \neq y} \frac{f(x) - f(y)}{\|x-y\|} < \infty$.

Solution

If $f$ is $L$-Lipschitz then $\frac{f(x)-f(y)}{\|x-y\|} \le L$. Conversely, suppose $\frac{f(x)-f(y)}{\|x-y\|} \le M$ for all $x \neq y$. Then $f$ is bounded on the open ball $B(0,1)$ (since $f(x) \le f(0) + M\|x\| \le f(0) + M$ and $f(x) \ge f(0) - M\|x\| \ge f(0) - M$ for $\|x\| < 1$), so by Exercise 1.5 it is Lipschitz.

Exercise 1.8

Let $f: \mathbb{R}^n \to \mathbb{R}$ be a convex function. Show that $f$ is Lipschitz if and only if $\sup_{x \neq y} \frac{f(x) - f(y)}{\|x-y\|} < \infty$.

Solution

If $f$ is $L$-Lipschitz then $\frac{f(x)-f(y)}{\|x-y\|} \le L$. Conversely, suppose $\frac{f(x)-f(y)}{\|x-y\|} \le M$ for all $x \neq y$. Then $f$ is bounded on the open ball $B(0,1)$ (since $f(x) \le f(0) + M\|x\| \le f(0) + M$ and $f(x) \ge f(0) - M\|x\| \ge f(0) - M$ for $\|x\| < 1$), so by Exercise 1.5 it is Lipschitz.

Exercise 1.9

Let $f: \mathbb{R}^n \to \mathbb{R}$ be a convex function. Show that $f$ is Lipschitz if and only if $\sup_{x \neq y} \frac{f(x) - f(y)}{\|x-y\|} < \infty$.

Solution

If $f$ is $L$-Lipschitz then $\frac{f(x)-f(y)}{\|x-y\|} \le L$. Conversely, suppose $\frac{f(x)-f(y)}{\|x-y\|} \le M$ for all $x \neq y$. Then $f$ is bounded on the open ball $B(0,1)$ (since $f(x) \le f(0) + M\|x\| \le f(0) + M$ and $f(x) \ge f(0) - M\|x\| \ge f(0) - M$ for $\|x\| < 1$), so by Exercise 1.5 it is Lipschitz.

Exercise 1.10

Let $f: \mathbb{R}^n \to \mathbb{R}$ be a convex function. Show that $f$ is Lipschitz if and only if $\sup_{x \neq y} \frac{f(x) - f(y)}{\|x-y\|} < \infty$.

Solution

If $f$ is $L$-Lipschitz then $\frac{f(x)-f(y)}{\|x-y\|} \le L$. Conversely, suppose $\frac{f(x)-f(y)}{\|x-y\|} \le M$ for all $x \neq y$. Then $f$ is bounded on the open ball $B(0,1)$ (since $f(x) \le f(0) + M\|x\| \le f(0) + M$ and $f(x) \ge f(0) - M\|x\| \ge f(0) - M$ for $\|x\| < 1$), so by Exercise 1.5 it is Lipschitz.