2024.05.30

Strong/weak Calabi-Yau/Calabi-Yau correspondence and D-brane transports

Kentaro Hori

Based on

KH, David Tong 2006

Manfred Herbst, KH, David Page 2008

KH 2011

KH, Mauricio Romo 2013

Richard Eager, KH, Johanna Knapp, Mauricio Romo 2024

Families of 2d (2,2) superconformal field theories can be obtained as the infra-red limits of simple gauge theories called

the **gauged linear sigma models**.


$G = U(1)$

$V = \mathbb{C}(1)^{\oplus 5} \oplus \mathbb{C}(-5)$

$x_1, \dots, x_5 \qquad p$

$W = p\, f(x_1, \dots, x_5)$

quintic polynomial



quintic Calabi-Yau 3-fold

$X_f = \{ f = 0 \} \subset \mathbb{CP}^4$

CY/LG correspondence


Witten 1993

Landau-Ginzburg orbifold

$(\mathbb{C}^5, f(x))/\mathbb{Z}_5$

$G = U(1)$

$V = \mathbb{C}(1)^{\oplus 2} \oplus \mathbb{C}(-2)$




Eguchi-Hanson space

$\mathcal{O}(-2)$


$\downarrow$

$\mathbb{CP}^1$



Orbifold


$\mathbb{C}^2/\{\pm 1\}$




McKay correspondence

$G = U(1)$


$V = \mathbb{C}(1)^{\oplus 2} \oplus \mathbb{C}(-1)^{\oplus 2}$




$\mathcal{O}(-1) \oplus \mathcal{O}(-1)$
$\downarrow$
$\mathbb{CP}^1$



conifold



flop



$\mathcal{O}(-1) \oplus \mathcal{O}(-1)$
$\downarrow$
$\mathbb{CP}^1$

CY/CY correspondence

Witten 1993

In these models, the phases are **weakly coupled**:

the gauge symmetry is completely broken or broken to a finite subgroup, and the classical analysis is enough to read off the low energy behavior.

There are interesting models with **strongly coupled phases**:

a continuous gauge symmetry remains unbroken, and you need a quantum analysis to find the low energy behavior.

Rødland model

H Tong 2006

$$G = U(2)$$


$$V = (\det^{-1})^{\oplus 7} \oplus (\mathbb{C}^2)^{\oplus 7} \quad \ni (p^1, \cdots, p^7, x_1, \cdots, x_7)$$

$$W = \sum_{i,j,k} A^{ij}_k \, p^k [x_i x_j]$$

$$[x_i x_j] := x_i^1 x_j^2 - x_i^2 x_j^2 \,,$$

write $$W = \sum_{ij} A^{ij}(p)[x_i x_j] = \sum_k p^k A_k(x)$$

$A^{ij}(p)$: mass matrix



$U(2) \to \{1\}$

"weakly coupled phase"

$\exists$ Coulomb branch


at $\begin{cases} \zeta_a = \mp \log\left(2\cos\left(\frac{\pi a}{7}\right)\right) \\ \theta_a \equiv \pi a \end{cases}$ $a = 1, 2, 3.$

$SU(2)$ unbroken

"strongly coupled phase"

$\xi \ll 0$: $SU(2)$ gauge theory fibered over $\mathbb{CP}^6 \ni p$:

seven fundamentals with mass matrix $A^{ij}(p) \xrightarrow{\text{rank}} \{4, 6\}$




$\operatorname{rank} A(p) = 6$: ~~SUSY~~

$\operatorname{rank} A(p) = 4$: 3 massless $\square$

$\xrightarrow{\text{I.R.}}$ free theory of composites

$[X_1 X_2]$, $[X_2 X_3]$, $[X_3 X_2]$

$\xrightarrow{\text{I.R.}}$ σ-model with target Y_A **Pfaffian variety**



$$X_A = \{ A_1(x) = \cdots = A_7(x) = 0 \} \subset G(2,7)$$

Grassmannian

$\exists$ Coulomb branch

at $\begin{cases} \zeta_a = 7 \log\left(2\cos\left(\frac{\pi a}{7}\right)\right) \\ \theta_a \equiv \pi a \end{cases}$ $a = 1, 2, 3.$

$$Y_A = \{ \operatorname{rk} A(p) = 4 \} \subset \mathbb{CP}^6$$

Pfaffian

This is an example of a

strong/weak Calabi-Yau/Calabi-Yau correspondence

known also as **Pfaffian/Grassmannian correspondence**

There are other examples:

Hosono-Takagi model

H 2011

$$G = \frac{O(2) \times U(1)}{\{(\pm 1_2, \pm 1)\}}$$


$$V = \mathbb{C}_+(-2)^{\oplus 5} \oplus \mathbb{C}^2(1)^{\oplus 5} \ni (p^1, \cdots, p^5, x_1, \cdots, x_5)$$

$$W = \sum_{i,j,k} S_k^{ij} p^k (x_i x_j)$$

$$(x_i x_j) := x_i^1 x_j^1 + x_i^2 x_j^2$$

write $W = \sum_{ij} S^{ij}(p)(x_i x_j) = \sum_k p^k S_k(x)$

mass matrix



$G \to \{1\}$

"weakly coupled phase"

$\exists$ Coulomb branch


at $\begin{cases} \zeta_a = 5 \log\left(2\cos\left(\frac{\pi a}{5}\right)\right) \\ \theta_a \equiv \pi a + \pi \end{cases}$ $a = 0, 1, 2.$

$O(2)$ unbroken

"strongly coupled phase"

$\zeta \ll 0$: $O(2)$ gauge theory fibered over $\mathbb{CP}^4 \ni p$:

five fundamentals with mass matrix $S^{ij}(p) \xrightarrow{\text{rank}} \{3, 4, 5\}$




$\text{rank}\, S(p) = 5$: ~~SUSY~~

$\text{rank}\, S(p) = 4$: two vacua

$\text{rank}\, S(p) = 3$: single vacuum

$\xrightarrow{\text{I.R.}}$ σ-model with target

ramified double cover $\tilde{Y}_S \to Y_S \cup C_S$



$$X_S = \{ S_1(x) = \cdots = S_5(x) = 0 \} \subset \frac{\mathbb{CP}^4 \times \mathbb{CP}^4}{\text{exchange}}$$

Reye congruence

$\exists$ Coulomb branch

at $\begin{cases} \zeta_a = 5 \log\left(2\cos\left(\frac{\pi a}{5}\right)\right) \\ \theta_a \equiv \pi a + \pi \end{cases}$ $a = 0, 1, 2.$

$$\widetilde{Y}_S \longrightarrow Y_S = \{ \operatorname{rk} S(p) \leq 4 \}$$
$$\cup$$
$$C_S = \{ \operatorname{rk} S(p) = 3 \}$$

There are dual models where

strong and **weak** are exchanged.

Constructed from 2d Seiberg duality. H 2011

Dual Rødland

H 2011


$$G^\vee = \frac{USp(4) \times U(1)}{\{(\pm \mathbb{1}_4, \pm 1)\}}$$

$$V^\vee = \mathbb{C}(-2)^{\oplus 7} \oplus \mathbb{C}^4(-1)^{\oplus 7} \oplus \mathbb{C}(2)^{\oplus \binom{7}{2}} \ni (p^k ; \tilde{x}^i ; a_{ij})$$

$$W^\vee = \sum_{i,j,k} A_k^{ij} p^k a_{ij} + \sum_{i,j} a_{ij} [\tilde{x}^i \tilde{x}^j]$$

$$[\tilde{x}^i \tilde{x}^j] = J^{ab} \tilde{x}_a^i \tilde{x}_b^j$$

$$a_{ij} = -a_{ji}$$



$US_p(4)$ unbroken


"strongly coupled phase"

$\exists$ Coulomb branch

at $\begin{cases} \zeta_a^\vee = 7 \log\left(2\cos\left(\frac{\pi a}{7}\right)\right) \\ \theta_a^\vee \equiv \pi a \end{cases}$ $a = 1, 2, 3.$

$G^\vee \to \{1\}$ (superpotential needed)

"weakly coupled phase"



$$\{\ rk\ a = 2\ ,\ \sum_{ij} A_k^{ij} a_{ij} = 0\ \} \subset \mathbb{CP}^{20}$$

$$a_{ij} = [x_i x_j] \qquad \cong X_A$$

Plücker

$\exists$ Coulomb branch

at $\begin{cases} \zeta_a^\vee = 7 \log\left(2\cos\left(\frac{\pi a}{7}\right)\right) \\ \theta_a^\vee \equiv \pi a \end{cases}$ $a = 1, 2, 3.$

$$\{\ A_{ij}(p) + [\tilde{x}^i \tilde{x}^j] = 0\ \}^S / G_C^\vee \cong Y_A$$

$$\Downarrow$$

$$rk\, A(p) = 4$$

Dual Hosono-Takagi

H 2011


$$G^\vee = \frac{SO(4)\times U(1)}{\{(\pm \mathbb{1}_4, \pm 1)\}}$$

$$V^\vee = \mathbb{C}(-2)^{\oplus 5} \oplus \mathbb{C}^4(-1)^{\oplus 5} \oplus \mathbb{C}(2)^{\oplus 15} \ni (p^k; \tilde{x}^i; s_{ij})$$

$$W^\vee = \sum_{i,j,k} S_k^{ij} p^k s_{ij} + \sum_{ij} s_{ij}(\tilde{x}^i \tilde{x}^j)$$

$$[\tilde{x}^i \tilde{x}^j] = \delta^{ab} \tilde{x}^i_a \tilde{x}^j_b$$

$$s_{ij} = s_{ji}$$



$SO(4)$ unbroken


"strongly coupled phase"

$\exists$ Coulomb branch

at $\begin{cases} \zeta_a^\vee = 5 \log\left(\cos\left(\frac{\pi a}{5}\right)\right) \\ \theta_a^\vee \equiv \pi a + \pi \end{cases}$ $a = 0, 1, 2.$

$G^\vee \to \{1\}$ (superpotential needed)

"weakly coupled phase"



$$\left\{ \operatorname{rank} S = 2,\ \sum_{ij} S^{ij} s_{ij} = 0 \right\} \subset \mathbb{CP}^{14}$$

$$s_{ij} = (x_i x_j) \qquad \cong X_S$$

$\exists$ Coulomb branch

at $\begin{cases} \xi_a^\vee = 5 \log\left(\cos\left(\frac{\pi a}{5}\right)\right) \\ \theta_a^\vee \equiv \pi a + \pi \end{cases}$ $a = 0, 1, 2.$

$$\left\{ S^{ij}(p) + (\tilde{x}^i \tilde{x}^j) = 0 \right\}^5 / G_C^\vee \cong \tilde{Y}_S$$

$$\downarrow \qquad\qquad \downarrow$$

$$\{ \operatorname{rk} S(p) \leq 4 \} = Y_S$$


Some observables are renormalization group invariant, and have special dependence on moduli, e.g. holomorphic.

Categories of **D-branes** are examples of such observables :

for B-branes, it is locally constant on Kahler moduli.

The category of B-branes in the sigma model with target X is the derived category of coherent sheaves $D^b(X)$.


Categories of B-branes



$$D^b(X_A)$$

$$D^b(Y_A)$$

A path



An equivalence of categories


$$D^b(X_A)$$

$$\cong$$

$$\downarrow$$

$$D^b(Y_A)$$

Another path



Another equivalence of categories


$$D^b(X_A)$$

$$\cong$$

$$\downarrow$$

$$D^b(Y_A)$$

Loops



Autoequivalences of categories

$D^b(X_A)$

$\simeq$

$\simeq$

$D^b(Y_A)$

What are these equivalences and monodromies?

By RG invariance, we may be able to study it using

the gauge theory realization of the theory.

In GLSM with gauge group G, matter V, superpotential W,

a classical B-brane is specified by

(M, Q) ··· a G *-equivariant matrix factorization of* (V, W)

$M = M^{ev} \oplus M^{od}$ a $\mathbb{Z}_2$ graded representation of G

$Q : V \longrightarrow End^{od}(M)$ a G-equivariant polynomial function

s.t. $Q^2 = W\, id_M$

+ an additional grading compatible with $\mathbb{Z}_2$.

They form a category $D_G(V, W)$.

However, a classical B-brane may induce instability in quantum theory, especially when a continuous gauge symmetry remains unbroken.

E.g. Around phase boundaries

Inside strongly coupled phases


Theories with simple gauge groups

We would like to find a condition on (M, Q) to define a well behaved B-brane in the quantum theory.

A proposal

Convergence of **hemisphere partition function**

HR, Sugishita-Terashima, Honda-Okuda 2013

$$\left\langle \text{} \right\rangle = \int_{\Upsilon} d^l\sigma \prod_{\alpha>0} \Gamma(\langle \alpha, \sigma \rangle) \sinh(\pi\langle \alpha, \sigma \rangle) \times \prod_i \Gamma\left(i\langle Q_i, \sigma \rangle + \frac{R_i}{2}\right) \times e^{i\langle t, \sigma \rangle} \cdot \mathrm{tr}_M\left(e^{\pi i R_M} e^{2\pi\sigma}\right)$$

$$V|_{T\times U(1)_V} = \bigoplus_i \mathbb{C}(Q_i, R_i)$$ weight decomposition

This imposes a severe constraint on representations of G that can be included in M.


For theories with Abelian gauge groups, this reproduces the **grade restriction rule** for D-brane transport across phase boundaries. HHP 2008

We shall continue to call the constraint **grade restriction rule**.

Example G simple, V symmetric: $\{\pm Q_i\}_{i>0} \cup \{\sigma's\}$

The representation of highest weight λ can be included in M when $\langle \lambda+\rho, \sigma\rangle < \frac{1}{2}\sum_{i>0}|\langle Q_i, \sigma\rangle| \quad \forall \sigma \neq 0 \in it.$

E.g. super-QCD $G = USp(k)$, $V = (\mathbb{C}^k)^{\oplus N}$ (N odd):

$S_Y(\mathbb{C}^k)^{\otimes|Y|}_0$ for Y in $\frac{k}{2}$ { 

Young symmetrizer J-traceless

Consistent with 2d Seiberg duality H2011.

Let us apply this to Rødland model, Hosono-Takagi model,

and their duals. Eager-H-Knapp-Romo 2014-2024

These models have gauge groups of the form $G = \dfrac{H \times U(1)}{\{(\pm 1_H, \pm 1)\}}$

with $H = USp(k), O(k), SO(k)$ e.g. $U(2) = \dfrac{USp(2) \times U(1)}{\{(\pm 1_2, \pm 1)\}}$.

Irreps of such groups is of the form $Y(i)$

- Y Young diagram specifying an irrep of $H \subset G$
- $U(1) \subset G$ acts by $\lambda \mapsto \lambda^{|Y|+2i}$

Grade restriction rule in strongly coupled phases:

Rodland $\zeta \ll 0$: $\mathbb{C}(i)$, $\square(i)$, $\square\square(i)$

Dual Rodland $\zeta^\vee \gg 0$ & $\zeta^\vee \ll 0$: $\mathbb{C}(i)$, $\square(i)$, $\begin{smallmatrix}\square\\\square\end{smallmatrix}(i)$

Hosono-Takagi $\zeta \ll 0$: $\mathbb{C}_\pm(i)$, $\square(i)$, $\square\square(i)$


Dual Hosono-Takagi $\zeta^\vee \gg 0$ & $\zeta^\vee \ll 0$: $\mathbb{C}(i)$, $\square(i)$, $\begin{smallmatrix}\square\\\square\end{smallmatrix}_{\pm}(i)$


$O(2)$ representation $\mathbb{C}_\pm$: $\begin{pmatrix}1 & 0\\ 0 & -1\end{pmatrix} \mapsto \pm 1$

$SO(4)$ representation $\begin{smallmatrix}\square\\\square\end{smallmatrix}_{\pm}$: $\begin{cases}\text{self-dual}\\ \text{anti-self-dual}\end{cases}$ 2-form

Grade restriction rule for D-brane transport

Rødland model:






$\mathbb{C}(-2)$ $\mathbb{C}(-1)$ $\mathbb{C}$ $\mathbb{C}(1)$ $\mathbb{C}(2)$ $\cdots$ $\mathbb{C}(5)$ $\mathbb{C}(6)$ $\mathbb{C}(7)$ $\mathbb{C}(8)$

$\square(-2)$ $\square(-1)$ $\square$ $\square(1)$ $\square(2)$ $\cdots$ $\square(5)$ $\square(6)$ $\square(7)$ $\square(8)$


$\square\square(-2)$ $\square\square(-1)$ $\square\square$ $\square\square(1)$ $\square\square(2)$ $\cdots$ $\square\square(5)$ $\square\square(6)$ $\square\square(7)$ $\square\square(8)$



$\mathbb{C}(-2)$ $\mathbb{C}(-1)$ $\mathbb{C}$ $\mathbb{C}(1)$ $\mathbb{C}(2)$ $\cdots$ $\mathbb{C}(5)$ $\mathbb{C}(6)$ $\mathbb{C}(7)$ $\mathbb{C}(8)$

$\square(-2)$ $\square(-1)$ $\square$ $\square(1)$ $\square(2)$ $\cdots$ $\square(5)$ $\square(6)$ $\square(7)$ $\square(8)$


$\square\square(-2)$ $\square\square(-1)$ $\square\square$ $\square\square(1)$ $\square\square(2)$ $\cdots$ $\square\square(5)$ $\square\square(6)$ $\square\square(7)$ $\square\square(8)$




$\mathbb{C}(-2)$ $\mathbb{C}(-1)$ $\mathbb{C}$ $\mathbb{C}(1)$ $\mathbb{C}(2)$ $\cdots$ $\mathbb{C}(5)$ $\mathbb{C}(6)$ $\mathbb{C}(7)$ $\mathbb{C}(8)$

$\square(-2)$ $\square(-1)$ $\square$ $\square(1)$ $\square(2)$ $\cdots$ $\square(5)$ $\square(6)$ $\square(7)$ $\square(8)$

$\square\square(-2)$ $\square\square(-1)$ $\square\square$ $\square\square(1)$ $\square\square(2)$ $\cdots$ $\square\square(5)$ $\square\square(6)$ $\square\square(7)$ $\square\square(8)$



$\mathbb{C}(-2)$	$\mathbb{C}(-1)$	$\mathbb{C}$	$\mathbb{C}(1)$	$\mathbb{C}(2)$	$\cdots$	$\mathbb{C}(5)$	$\mathbb{C}(6)$	$\mathbb{C}(7)$	$\mathbb{C}(8)$
$\square(-2)$	$\square(-1)$	$\square$	$\square(1)$	$\square(2)$	$\cdots$	$\square(5)$	$\square(6)$	$\square(7)$	$\square(8)$
$\square\square(-2)$	$\square\square(-1)$	$\square\square$	$\square\square(1)$	$\square\square(2)$	$\cdots$	$\square\square(5)$	$\square\square(6)$	$\square\square(7)$	$\square\square(8)$




$\mathbb{C}(-2)$ $\mathbb{C}(-1)$ $\mathbb{C}$ $\mathbb{C}(1)$ $\mathbb{C}(2)$ $\cdots$ $\mathbb{C}(5)$ $\mathbb{C}(6)$ $\mathbb{C}(7)$ $\mathbb{C}(8)$

$\square(-2)$ $\square(-1)$ $\square$ $\square(1)$ $\square(2)$ $\cdots$ $\square(5)$ $\square(6)$ $\square(7)$ $\square(8)$

$\square\square(-2)$ $\square\square(-1)$ $\square\square$ $\square\square(1)$ $\square\square(2)$ $\cdots$ $\square\square(5)$ $\square\square(6)$ $\square\square(7)$ $\square\square(8)$

For each homotopy class of paths (“window”) **w**

between $\zeta \gg 0$ and $\zeta \ll 0$, such as




let $\mathcal{T}_{\mathbf{w}}$ be the subcategory of


equivariant matrix factorizations made of

grade restricted representations with respect to **w**.

S : the one at $\zeta \ll 0$.


Then, we can make equivalences






c.f. Addington-Donovan-Segal 2014

Monodromy







$\mathbb{C}(-2)$ $\mathbb{C}(-1)$ $\mathbb{C}$ $\mathbb{C}(1)$ $\mathbb{C}(2)$ $\cdots$ $\mathbb{C}(5)$ $\mathbb{C}(6)$ $\mathbb{C}(7)$ $\mathbb{C}(8)$

$\square(-2)$ $\square(-1)$ $\square$ $\square(1)$ $\square(2)$ $\cdots$ $\square(5)$ $\square(6)$ $\square(7)$ $\square(8)$

$\square\square(-2)$ $\square\square(-1)$ $\square\square$ $\square\square(1)$ $\square\square(2)$ $\cdots$ $\square\square(5)$ $\square\square(6)$ $\square\square(7)$ $\square\square(8)$


W
W
$MF_{U(2)}(W)$
U
S
U
U
$\mathcal{J}_W$
$\mathcal{J}_W$
W
W
$D^b(Y_A)$
$D^b(X_A)$



$\mathbb{C}(-2)$ $\mathbb{C}(-1)$ $\mathbb{C}$ $\mathbb{C}(1)$ $\mathbb{C}(2)$ $\cdots$ $\mathbb{C}(5)$ $\mathbb{C}(6)$ $\mathbb{C}(7)$ $\mathbb{C}(8)$

$\square(-2)$ $\square(-1)$ $\square$ $\square(1)$ $\square(2)$ $\cdots$ $\square(5)$ $\square(6)$ $\square(7)$ $\square(8)$

$\square\square(-2)$ $\square\square(-1)$ $\square\square$ $\square\square(1)$ $\square\square(2)$ $\cdots$ $\square\square(5)$ $\square\square(6)$ $\square\square(7)$ $\square\square(8)$

Want to replace $\mathbb{C}$ by $\mathbb{C}^{(7)}$ in the phase $\zeta \ll 0$.

—— Can be done with the brane $\mathcal{K}$


$$\mathbb{C} \underset{p}{\overset{A(x)}{\rightleftarrows}} \mathbb{C}^{7}(1) \underset{p}{\overset{A(x)}{\rightleftarrows}} \mathbb{C}^{\binom{7}{2}}(2) \underset{p}{\overset{A(x)}{\rightleftarrows}} \mathbb{C}^{\binom{7}{3}}(3) \underset{p}{\overset{A(x)}{\rightleftarrows}} \mathbb{C}^{\binom{7}{4}}(4) \underset{p}{\overset{A(x)}{\rightleftarrows}} \mathbb{C}^{\binom{7}{5}}(5) \underset{p}{\overset{A(x)}{\rightleftarrows}} \mathbb{C}^{\binom{7}{6}}(6) \underset{p}{\overset{A(x)}{\rightleftarrows}} \mathbb{C}(7)$$

0 1 2 3 4 5 6 7

$$Q = \sum_{k=1}^{7} \left(p^k \eta_k + A_k(x) \bar{\eta}^k \right) \qquad \{ \eta_k, \bar{\eta}^\ell \} = \delta_k^\ell$$

which is empty, $\mathcal{K} \longmapsto 0$, as

$$\{ Q, Q^+ \} = \| p \|^2 + \| A(x) \|^2 > 0 \quad \text{if} \quad \zeta \ll 0.$$




$\mathbb{C}(-2)$ $\mathbb{C}(-1)$ $\mathbb{C}$ $\mathbb{C}(1)$ $\mathbb{C}(2)$ $\cdots$ $\mathbb{C}(5)$ $\mathbb{C}(6)$ $\mathbb{C}(7)$ $\mathbb{C}(8)$

$\square(-2)$ $\square(-1)$ $\square$ $\square(1)$ $\square(2)$ $\cdots$ $\square(5)$ $\square(6)$ $\square(7)$ $\square(8)$


$\square\square(-2)$ $\square\square(-1)$ $\square\square$ $\square\square(1)$ $\square\square(2)$ $\cdots$ $\square\square(5)$ $\square\square(6)$ $\square\square(7)$ $\square\square(8)$

Replacement $\square \longrightarrow \square(7)$ in the phase $\zeta \ll 0$

can be done with $\square \otimes K$:



which is empty there.




$\mathbb{C}(-2)$ $\mathbb{C}(-1)$ $\mathbb{C}$ $\mathbb{C}(1)$ $\mathbb{C}(2)$ $\cdots$ $\mathbb{C}(5)$ $\mathbb{C}(6)$ $\mathbb{C}(7)$ $\mathbb{C}(8)$

$\square(-2)$ $\square(-1)$ $\square$ $\square(1)$ $\square(2)$ $\cdots$ $\square(5)$ $\square(6)$ $\square(7)$ $\square(8)$

$\square\square(-2)$ $\square\square(-1)$ $\square\square$ $\square\square(1)$ $\square\square(2)$ $\cdots$ $\square\square(5)$ $\square\square(6)$ $\square\square(7)$ $\square\square(8)$

Replacement $\square\square(-1) \to \square\square(6)$ in the phase $\zeta \ll 0$

can be done with $\square\square \otimes K(-1)$:




which is empty there.

In the phase $\zeta \gg 0$,


GLSM		non-linear σ-model	
$\mathcal{K}$	$\longmapsto$	$\mathcal{O}$	the structure sheaf of X_A
$\square \otimes \mathcal{K}$	$\longmapsto$	$S^\vee$	a rank 2 vector bundle
$\square\square \otimes \mathcal{K}(-1)$	$\longmapsto$	$\mathrm{Sym}^2 S^\vee(-1)$	rank 3

under the RG flow.

Seidel-Thomas twists by




Dual Rødland model



$\cdots \mathbb{C}(-2)\ \ \mathbb{C}(-1)\ \ \mathbb{C}\ \ \mathbb{C}(1)\ \ \mathbb{C}(2)\ \ \mathbb{C}(3)\ \cdots\ \mathbb{C}(19)\ \ \mathbb{C}(20)\ \ \mathbb{C}(21)\ \ \mathbb{C}(22)\ \cdots$

$\cdots \square(-2)\ \ \square(-1)\ \ \square\ \ \square(1)\ \ \square(2)\ \ \square(3)\ \cdots\ \square(19)\ \ \square(20)\ \ \square(21)\ \ \square(22)\ \cdots$


$\cdots \boxminus(-2)\ \ \boxminus(-1)\ \ \boxminus\ \ \boxminus(1)\ \ \boxminus(2)\ \ \boxminus(3)\ \cdots\ \boxminus(19)\ \ \boxminus(20)\ \ \boxminus(21)\ \ \boxminus(22)\ \cdots$



$\cdots \mathbb{C}(-2)\ \mathbb{C}(-1)\ \mathbb{C}\ \mathbb{C}(1)\ \mathbb{C}(2)\ \mathbb{C}(3)\ \cdots\cdots\ \mathbb{C}(19)\ \mathbb{C}(20)\ \mathbb{C}(21)\ \mathbb{C}(22)\ \cdots$

$\cdots \square(-2)\ \square(-1)\ \square\ \square(1)\ \square(2)\ \square(3)\ \cdots\cdots\ \square(19)\ \square(20)\ \square(21)\ \square(22)\ \cdots$


$\cdots \begin{smallmatrix}\square\\\square\end{smallmatrix}(-2)\ \begin{smallmatrix}\square\\\square\end{smallmatrix}(-1)\ \begin{smallmatrix}\square\\\square\end{smallmatrix}\ \begin{smallmatrix}\square\\\square\end{smallmatrix}(1)\ \begin{smallmatrix}\square\\\square\end{smallmatrix}(2)\ \begin{smallmatrix}\square\\\square\end{smallmatrix}(3)\ \cdots\cdots\ \begin{smallmatrix}\square\\\square\end{smallmatrix}(19)\ \begin{smallmatrix}\square\\\square\end{smallmatrix}(20)\ \begin{smallmatrix}\square\\\square\end{smallmatrix}(21)\ \begin{smallmatrix}\square\\\square\end{smallmatrix}(22)\ \cdots$



$\cdots \mathbb{C}(-2)\ \ \mathbb{C}(-1)\ \ \mathbb{C}\ \ \mathbb{C}(1)\ \ \mathbb{C}(2)\ \ \mathbb{C}(3)\ \cdots\ \mathbb{C}(19)\ \ \mathbb{C}(20)\ \ \mathbb{C}(21)\ \ \mathbb{C}(22)\ \cdots$

$\cdots \square(-2)\ \ \square(-1)\ \ \square\ \ \square(1)\ \ \square(2)\ \ \square(3)\ \cdots\ \square(19)\ \ \square(20)\ \ \square(21)\ \ \square(22)\cdots$


$\cdots \begin{smallmatrix}\square\\\square\end{smallmatrix}(-2)\ \ \begin{smallmatrix}\square\\\square\end{smallmatrix}(-1)\ \ \begin{smallmatrix}\square\\\square\end{smallmatrix}\ \ \begin{smallmatrix}\square\\\square\end{smallmatrix}(1)\ \ \begin{smallmatrix}\square\\\square\end{smallmatrix}(2)\ \ \begin{smallmatrix}\square\\\square\end{smallmatrix}(3)\ \cdots\ \begin{smallmatrix}\square\\\square\end{smallmatrix}(19)\ \ \begin{smallmatrix}\square\\\square\end{smallmatrix}(20)\ \ \begin{smallmatrix}\square\\\square\end{smallmatrix}(21)\ \ \begin{smallmatrix}\square\\\square\end{smallmatrix}(22)\cdots$




$\cdots \mathbb{C}(-2)\ \ \mathbb{C}(-1)\ \ \mathbb{C}\ \ \mathbb{C}(1)\ \ \mathbb{C}(2)\ \ \mathbb{C}(3)\ \cdots\ \mathbb{C}(19)\ \ \mathbb{C}(20)\ \ \mathbb{C}(21)\ \ \mathbb{C}(22)\ \cdots$


$\cdots \square(-2)\ \ \square(-1)\ \ \square\ \ \square(1)\ \ \square(2)\ \ \square(3)\ \cdots\ \square(19)\ \ \square(20)\ \ \square(21)\ \ \square(22)\ \cdots$

$\cdots \begin{smallmatrix}\square\\\square\end{smallmatrix}(-2)\ \ \begin{smallmatrix}\square\\\square\end{smallmatrix}(-1)\ \ \begin{smallmatrix}\square\\\square\end{smallmatrix}\ \ \begin{smallmatrix}\square\\\square\end{smallmatrix}(1)\ \ \begin{smallmatrix}\square\\\square\end{smallmatrix}(2)\ \ \begin{smallmatrix}\square\\\square\end{smallmatrix}(3)\ \cdots\ \begin{smallmatrix}\square\\\square\end{smallmatrix}(19)\ \ \begin{smallmatrix}\square\\\square\end{smallmatrix}(20)\ \ \begin{smallmatrix}\square\\\square\end{smallmatrix}(21)\ \ \begin{smallmatrix}\square\\\square\end{smallmatrix}(22)\cdots$


$\zeta^\vee$
$\theta^\vee$
-17π


$\mathbb{C}(-2)$ $\mathbb{C}(-1)$ $\mathbb{C}$ $\mathbb{C}(1)$ $\mathbb{C}(2)$ $\mathbb{C}(3)$ $\mathbb{C}(19)$ $\mathbb{C}(20)$ $\mathbb{C}(21)$ $\mathbb{C}(22)$
$\square(-2)$ $\square(-1)$ $\square$ $\square(1)$ $\square(2)$ $\square(3)$ $\square(19)$ $\square(20)$ $\square(21)$ $\square(22)$
$\square\!\square(-2)$ $\square\!\square(-1)$ $\square\!\square$ $\square\!\square(1)$ $\square\!\square(2)$ $\square\!\square(3)$ $\square\!\square(19)$ $\square\!\square(20)$ $\square\!\square(21)$ $\square\!\square(22)$

For each window **W** such as





$$D_{G^\vee}(V^\vee, W^\vee)$$


$$\bigcup_{S^\vee} \bigcup_{J_{\mathbf{W}}}$$


$$D^b(Y_A) \cong \qquad \cong D^b(X_A)$$


Monodromy




$\zeta^\vee$
$\theta^\vee$
-17π
$\cdots \mathbb{C}(-2)$ $\mathbb{C}(-1)$ $\mathbb{C}$ $\mathbb{C}(1)$ $\mathbb{C}(2)$ $\mathbb{C}(3)$ $\cdots$ $\mathbb{C}(19)$ $\mathbb{C}(20)$ $\mathbb{C}(21)$ $\mathbb{C}(22)$ $\cdots$
$\cdots \square(-2)$ $\square(-1)$ $\square$ $\square(1)$ $\square(2)$ $\square(3)$ $\cdots$ $\square(19)$ $\square(20)$ $\square(21)$ $\square(22)$ $\cdots$
$\cdots \square\!\square(-2)$ $\square\!\square(-1)$ $\square\!\square$ $\square\!\square(1)$ $\square\!\square(2)$ $\square\!\square(3)$ $\cdots$ $\square\!\square(19)$ $\square\!\square(20)$ $\square\!\square(21)$ $\square\!\square(22)$ $\cdots$


$\zeta^\vee$
$\theta^\vee$
-17π
$\cdots \mathbb{C}(-2)\ \mathbb{C}(-1)\ \mathbb{C}\ \mathbb{C}(1)\ \mathbb{C}(2)\ \mathbb{C}(3)\ \cdots\cdots\ \mathbb{C}(19)\ \mathbb{C}(20)\ \mathbb{C}(21)\ \mathbb{C}(22)\ \cdots$
$\cdots \square(-2)\ \square(-1)\ \square\ \square(1)\ \square(2)\ \square(3)\ \cdots\cdots\ \square(19)\ \square(20)\ \square(21)\ \square(22)\ \cdots$
$\cdots \square\!\square(-2)\ \square\!\square(-1)\ \square\!\square\ \square\!\square(1)\ \square\!\square(2)\ \square\!\square(3)\ \cdots\cdots\ \square\!\square(19)\ \square\!\square(20)\ \square\!\square(21)\ \square\!\square(22)\ \cdots$


$\zeta^\vee$
$\theta^\vee$
-17π


$\cdots \mathbb{C}(-2)$ $\mathbb{C}(-1)$ $\mathbb{C}$ $\mathbb{C}(1)$ $\mathbb{C}(2)$ $\mathbb{C}(3)$ $\cdots\cdots$ $\mathbb{C}(19)$ $\mathbb{C}(20)$ $\mathbb{C}(21)$ $\mathbb{C}(22)\cdots$
$\cdots \square(-2)$ $\square(-1)$ $\square$ $\square(1)$ $\square(2)$ $\square(3)$ $\cdots\cdots$ $\square(19)$ $\square(20)$ $\square(21)$ $\square(22)\cdots$
$\cdots \boxminus(-2)$ $\boxminus(-1)$ $\boxminus$ $\boxminus(1)$ $\boxminus(2)$ $\boxminus(3)$ $\cdots\cdots$ $\boxminus(19)$ $\boxminus(20)$ $\boxminus(21)$ $\boxminus(22)\cdots$

The replacement in the phase $\zeta^\vee \gg 0$ can be done

with the branes $K^\vee$, $\square \otimes K^\vee$, $\begin{array}{|c|}\hline \\ \hline \\ \hline \end{array} \otimes K^\vee(-1)$

where $K^\vee$ is

$$\mathbb{C} \underset{A(p)+[\tilde{x}\tilde{x}]}{\overset{a}{\rightleftarrows}} \mathbb{C}(1)^{\oplus 21} \underset{A(p)+[\tilde{x}\tilde{x}]}{\overset{a}{\rightleftarrows}} \mathbb{C}(2)^{\oplus \binom{21}{2}} \underset{A(p)+[\tilde{x}\tilde{x}]}{\overset{a}{\rightleftarrows}} \cdots \underset{A(p)+[\tilde{x}\tilde{x}]}{\overset{a}{\rightleftarrows}} \mathbb{C}(20)^{\oplus \binom{21}{3}} \underset{A(p)+[\tilde{x}\tilde{x}]}{\overset{a}{\rightleftarrows}} \mathbb{C}(21)$$

$$Q = \sum_{i,j} \left(a_{ij} \eta^{ij} + \left(A^{ij}(p) + [\tilde{x}^i \tilde{x}^j] \right) \bar{\eta}_{ij} \right) \; ; \; \{\eta_{ij}, \bar{\eta}^{kl}\} = \epsilon_{ij}^{kl}$$

which are empty, $K^\vee \longmapsto 0$, as

$$\{Q, Q^\dagger\} = \|a\|^2 + \|A(p) + [\tilde{x}\tilde{x}]\|^2 > 0 \quad \text{if} \quad \zeta^\vee \gg 0.$$

In the phase $\zeta^\vee \ll 0$

GLSM non-linear σ-model


$K^\vee \longmapsto \mathcal{O}$ the structure sheaf of Y_A

$\square \otimes K^\vee \longmapsto \mathcal{R}$ a rank 4 vector bundle


$\begin{array}{|c|}\hline \\ \hline \\ \hline \end{array} \otimes K^\vee(-1) \longmapsto \Lambda^2_0 \mathcal{R}(1)$ rank 5

under the RG flow.

Seidel-Thomas twists by




Hosono-Takagi model




$\mathbb{C}_{\pm}(-2)$ $\mathbb{C}_{\pm}(-1)$ $\mathbb{C}_{\pm}$ $\mathbb{C}_{\pm}(1)$ $\mathbb{C}_{\pm}(2)$ $\mathbb{C}_{\pm}(3)$ $\mathbb{C}_{\pm}(4)$ $\mathbb{C}_{\pm}(5)$ $\mathbb{C}_{\pm}(6)$

$\square(-2)$ $\square(-1)$ $\square$ $\square(1)$ $\square(2)$ $\square(3)$ $\square(4)$ $\square(5)$ $\square(6)$

$\square\square(-2)$ $\square\square(-1)$ $\square\square$ $\square\square(1)$ $\square\square(2)$ $\square\square(3)$ $\square\square(4)$ $\square\square(5)$ $\square\square(6)$




$\mathbb{C}_{\pm}(-2)$	$\mathbb{C}_{\pm}(-1)$	$\mathbb{C}_{\pm}$	$\mathbb{C}_{\pm}(1)$	$\mathbb{C}_{\pm}(2)$	$\mathbb{C}_{\pm}(3)$	$\mathbb{C}_{\pm}(4)$	$\mathbb{C}_{\pm}(5)$	$\mathbb{C}_{\pm}(6)$
□(-2)	□(-1)	□	□(1)	□(2)	□(3)	□(4)	□(5)	□(6)
□□(-2)	□□(-1)	□□	□□(1)	□□(2)	□□(3)	□□(4)	□□(5)	□□(6)



$\mathbb{C}_{\pm}(-2)$ $\mathbb{C}_{\pm}(-1)$ $\mathbb{C}_{\pm}$ $\mathbb{C}_{\pm}(1)$ $\mathbb{C}_{\pm}(2)$ $\mathbb{C}_{\pm}(3)$ $\mathbb{C}_{\pm}(4)$ $\mathbb{C}_{\pm}(5)$ $\mathbb{C}_{\pm}(6)$

$\square(-2)$ $\square(-1)$ $\square$ $\square(1)$ $\square(2)$ $\square(3)$ $\square(4)$ $\square(5)$ $\square(6)$


$\square\square(-2)$ $\square\square(-1)$ $\square\square$ $\square\square(1)$ $\square\square(2)$ $\square\square(3)$ $\square\square(4)$ $\square\square(5)$ $\square\square(6)$



$\mathbb{C}_{\pm}(-2)$ $\mathbb{C}_{\pm}(-1)$ $\mathbb{C}_{\pm}$ $\mathbb{C}_{\pm}(1)$ $\mathbb{C}_{\pm}(2)$ $\mathbb{C}_{\pm}(3)$ $\mathbb{C}_{\pm}(4)$ $\mathbb{C}_{\pm}(5)$ $\mathbb{C}_{\pm}(6)$


$\square(-2)$ $\square(-1)$ $\square$ $\square(1)$ $\square(2)$ $\square(3)$ $\square(4)$ $\square(5)$ $\square(6)$


$\square\square(-2)$ $\square\square(-1)$ $\square\square$ $\square\square(1)$ $\square\square(2)$ $\square\square(3)$ $\square\square(4)$ $\square\square(5)$ $\square\square(6)$



$\mathbb{C}_{\pm}(-2)$	$\mathbb{C}_{\pm}(-1)$	$\mathbb{C}_{\pm}$	$\mathbb{C}_{\pm}(1)$	$\mathbb{C}_{\pm}(2)$	$\mathbb{C}_{\pm}(3)$	$\mathbb{C}_{\pm}(4)$	$\mathbb{C}_{\pm}(5)$	$\mathbb{C}_{\pm}(6)$
□(-2)	□(-1)	□	□(1)	□(2)	□(3)	□(4)	□(5)	□(6)
□□(-2)	□□(-1)	□□	□□(1)	□□(2)	□□(3)	□□(4)	□□(5)	□□(6)

Then, we can make equivalences






c.f. Rennemo 2015

Monodromy

$\mathcal{E} = [\mathcal{O}(1,0) \oplus \mathcal{O}(0,1)]$


rank 2



rank 2

$\mathrm{Sym}^2_{\mathcal{O}} \mathcal{E}(-1)$

Dual Hosono-Takagi




$$\cdots \mathbb{C}(-2)\ \ \mathbb{C}(-1)\ \ \mathbb{C}\ \ \mathbb{C}(1)\ \ \mathbb{C}(2)\ \ \mathbb{C}(3)\ \cdots\ \mathbb{C}(13)\ \ \mathbb{C}(14)\ \ \mathbb{C}(15)\ \ \mathbb{C}(16)\ \cdots$$

$$\cdots \square(-2)\ \ \square(-1)\ \ \square\ \ \square(1)\ \ \square(2)\ \ \square(3)\ \cdots\ \square(13)\ \ \square(14)\ \ \square(15)\ \ \square(16)\ \cdots$$

$$\cdots \boxminus_+(-2)\ \ \boxminus_+(-1)\ \ \boxminus_+\ \ \boxminus_+(1)\ \ \boxminus_+(2)\ \ \boxminus_+(3)\ \cdots\ \boxminus_+(13)\ \ \boxminus_+(14)\ \ \boxminus_+(15)\ \ \boxminus_+(16)\ \cdots$$

$$\cdots \boxminus_-(-2)\ \ \boxminus_-(-1)\ \ \boxminus_-\ \ \boxminus_-(1)\ \ \boxminus_-(2)\ \ \boxminus_-(3)\ \cdots\ \boxminus_-(13)\ \ \boxminus_-(14)\ \ \boxminus_-(15)\ \ \boxminus_-(16)\ \cdots$$




$\cdots \mathbb{C}(-2)\ \mathbb{C}(-1)\ \mathbb{C}\ \mathbb{C}(1)\ \mathbb{C}(2)\ \mathbb{C}(3)\ \cdots\cdots\ \mathbb{C}(13)\ \mathbb{C}(14)\ \mathbb{C}(15)\ \mathbb{C}(16)\ \cdots$

$\cdots \square(-2)\ \square(-1)\ \square\ \square(1)\ \square(2)\ \square(3)\ \cdots\cdots\ \square(13)\ \square(14)\ \square(15)\ \square(16)\ \cdots$

$\cdots \boxminus_+(-2)\ \boxminus_+(-1)\ \boxminus_+\ \boxminus_+(1)\ \boxminus_+(2)\ \boxminus_+(3)\ \cdots\cdots\ \boxminus_+(13)\ \boxminus_+(14)\ \boxminus_+(15)\ \boxminus_+(16)\ \cdots$

$\cdots \boxminus_-(-2)\ \boxminus_-(-1)\ \boxminus_-\ \boxminus_-(1)\ \boxminus_-(2)\ \boxminus_-(3)\ \cdots\cdots\ \boxminus_-(13)\ \boxminus_-(14)\ \boxminus_-(15)\ \boxminus_-(16)\ \cdots$




$$\cdots \mathbb{C}(-2)\ \ \mathbb{C}(-1)\ \ \mathbb{C}\ \ \mathbb{C}(1)\ \ \mathbb{C}(2)\ \ \mathbb{C}(3)\ \cdots\ \mathbb{C}(13)\ \ \mathbb{C}(14)\ \ \mathbb{C}(15)\ \ \mathbb{C}(16)\ \cdots$$

$$\cdots \square(-2)\ \ \square(-1)\ \ \square\ \ \square(1)\ \ \square(2)\ \ \square(3)\ \cdots\ \square(13)\ \ \square(14)\ \ \square(15)\ \ \square(16)\ \cdots$$

$$\cdots \boxminus_+(-2)\ \ \boxminus_+(-1)\ \ \boxminus_+\ \ \boxminus_+(1)\ \ \boxminus_+(2)\ \ \boxminus_+(3)\ \cdots\ \boxminus_+(13)\ \ \boxminus_+(14)\ \ \boxminus_+(15)\ \ \boxminus_+(16)\ \cdots$$

$$\cdots \boxminus_-(-2)\ \ \boxminus_-(-1)\ \ \boxminus_-\ \ \boxminus_-(1)\ \ \boxminus_-(2)\ \ \boxminus_-(3)\ \cdots\ \boxminus_-(13)\ \ \boxminus_-(14)\ \ \boxminus_-(15)\ \ \boxminus_-(16)\ \cdots$$




$$\cdots \mathbb{C}(-2)\ \ \mathbb{C}(-1)\ \ \mathbb{C}\ \ \mathbb{C}(1)\ \ \mathbb{C}(2)\ \ \mathbb{C}(3)\ \cdots\ \mathbb{C}(13)\ \ \mathbb{C}(14)\ \ \mathbb{C}(15)\ \ \mathbb{C}(16)\ \cdots$$

$$\cdots \square(-2)\ \ \square(-1)\ \ \square\ \ \square(1)\ \ \square(2)\ \ \square(3)\ \cdots\ \square(13)\ \ \square(14)\ \ \square(15)\ \ \square(16)\ \cdots$$

$$\cdots \boxminus_+(-2)\ \ \boxminus_+(-1)\ \ \boxminus_+\ \ \boxminus_+(1)\ \ \boxminus_+(2)\ \ \boxminus_+(3)\ \cdots\ \boxminus_+(13)\ \ \boxminus_+(14)\ \ \boxminus_+(15)\ \ \boxminus_+(16)\ \cdots$$

$$\cdots \boxminus_-(-2)\ \ \boxminus_-(-1)\ \ \boxminus_-\ \ \boxminus_-(1)\ \ \boxminus_-(2)\ \ \boxminus_-(3)\ \cdots\ \boxminus_-(13)\ \ \boxminus_-(14)\ \ \boxminus_-(15)\ \ \boxminus_-(16)\ \cdots$$




$$\cdots \mathbb{C}(-2)\ \ \mathbb{C}(-1)\ \ \mathbb{C}\ \ \mathbb{C}(1)\ \ \mathbb{C}(2)\ \ \mathbb{C}(3)\ \cdots\ \mathbb{C}(13)\ \ \mathbb{C}(14)\ \ \mathbb{C}(15)\ \ \mathbb{C}(16)\cdots$$

$$\cdots \square(-2)\ \ \square(-1)\ \ \square\ \ \square(1)\ \ \square(2)\ \ \square(3)\ \cdots\ \square(13)\ \ \square(14)\ \ \square(15)\ \ \square(16)\cdots$$

$$\cdots \boxminus_+(-2)\ \ \boxminus_+(-1)\ \ \boxminus_+\ \ \boxminus_+(1)\ \ \boxminus_+(2)\ \ \boxminus_+(3)\ \cdots\ \boxminus_+(13)\ \ \boxminus_+(14)\ \ \boxminus_+(15)\ \ \boxminus_+(16)\cdots$$

$$\cdots \boxminus_-(-2)\ \ \boxminus_-(-1)\ \ \boxminus_-\ \ \boxminus_-(1)\ \ \boxminus_-(2)\ \ \boxminus_-(3)\ \cdots\ \boxminus_-(13)\ \ \boxminus_-(14)\ \ \boxminus_-(15)\ \ \boxminus_-(16)\cdots$$

For each window **W** such as



$$D_{G^\vee}(V^\vee, W^\vee)$$

$$\cup$$

$$S^\vee$$

$$\cup$$

$$\widetilde{J}^\vee_{\mathbf{W}}$$

$$\cong \qquad\qquad \cong$$

$$D^b(\widetilde{Y}_S) \qquad\qquad D^b(X_S)$$

Monodromy

$\Lambda^2_+ \mathcal{F}(1) \circ \Lambda^2_- \mathcal{F}$ rank 3

rank 4

