Non-Invertible Peccei-Quinn Symmetry and the Massless Quark Solution to the Strong CP Problem

Sungwoo Hong

KAIST

2211.07639: C Córdova, **SH**, S Koren, K Ohmori
2306.00912: T.D Brennan, **SH**
2309.05636: C Córdova, **SH**, LT Wang
2402.12453: C Córdova, **SH**, S Koren

The 12th KIAS Workshop on Particle Physics and Cosmology
2024 Korea-France STAR Workshop

Opening Remarks

Opening Remarks

Naturalness Problems and Global Symmetries

1. Electroweak Hierarch Problem

$$\left(\frac{\text{Gravity}}{\text{weak}}\right) \sim \left(\frac{v}{M_{pl}}\right)^2 \sim \left(\frac{100\ GeV}{10^{19}\ GeV}\right)^2 \sim 10^{-34} \ll 1$$

A source of challenge: **no apparent symmetry** acting on (generic) **scalar Φ**

Exception-1) Shift symmetry: Higgs = PNGB $\Rightarrow$ Composite Higgs / Little Higgs

Exception-2) Chiral symmetry (scalar $\leftrightarrow$ fermion): SUSY $\Rightarrow$ (N)MSSM

In these cases, hierarchy problem becomes **Technical Naturalness Problem.**

Opening Remarks

Naturalness Problems and Global Symmetries

2. Strong CP Problem

$$\tilde{J} = \mathbf{Im\,det}[y_u^\dagger y_u, y_d^\dagger y_d\,] \propto \sin\delta_{CKM} \sim O(1) \quad \textbf{vs} \quad \bar{\theta} \equiv \arg e^{-i\theta}\det(y_u y_d) \ll 1$$

"Jarlskog invariant"

source of challenge 1: **no clean symmetry structure**

CP (=T), Anomalous $U(1)_{PQ}$, flavor symmetry, …
renormalization of $\bar{\theta}$ from other CPV sources


source of challenge 2: the limit $\bar{\theta} \to 0$ does not enhance the symmetry of QFT

Strong CP problem = Dirac Naturalness Problem

Opening Remarks

Naturalness Problems and Global Symmetries

3. Flavor Problem [e.g. m_ν]



https://www.researchgate.net/figure/Mass-spectrum-of-standard-model-fermions-Charged-leptons-up-type-quarks-and-downtype_fig1_361578459

$M_\nu \sim 10^{-2}\ eV$

Requires Dynamical Explanation!


Several attractive theories exist.

(1) Seesaw models based on $U(1)_L$
(2) Extradimension, clockwork: localization
(3) …

Opening Remarks

Naturalness Problems and Global Symmetries

3. Flavor Problem [e.g. m_ν]



https://www.researchgate.net/figure/Mass-spectrum-of-standard-model-fermions-Charged-leptons-up-type-quarks-and-downtype_fig1_361578459

$M_\nu \sim 10^{-2}\ eV$


Requires Dynamical Explanation!

A source of challenge: ultimate mechanism still to be confirmed.

=> more feasible, testable, and motivating **theoretical ideas** should be laid out.


Opening Remarks

Generalized Global Symmetries



Opening Remarks

Generalized Global Symmetries



Opening Remarks

Generalized Global Symmetries in Physics

Well-motivated and timely to think about new ideas and breakthrough Generalized Global Symmetry can provide.

Opening Remarks

Generalized Global Symmetries in Particle Physics

0. Noninvertible Chiral Symmetry and Exponential Hierarchies '22 (C. Cordova, K. Ohmori)
   Noninvertible Global Symmetries in the Standard Model '22 (Y. Choi, H.T. Lam, S.-H Shao)

1. Neutrino Masses from Generalized Symmetry Breaking '22 (C. Cordova, **SH**, S. Koren, K. Ohmori)

2. Higher Flavor Symmetries in the Standard Model '22 (C. Cordova, S. Koren)

3. Coupling a Cosmic String to a TQFT '23 (T.D. Brennan, **SH**, LT Wang)
   Quantization of Axion-Gauge Couplings and Non-Invertible Higher Symmetries '23 (Y. Choi, M. Forslund, H. T. Lam, S-H. Shao)
   Axion-Gauge Coupling Quantization with a Twist '23 (M. Reece)
   Axion Domain Walls, Small Instantons, and Non-Invertible Symmetry Breaking '23 (C. Cordova, **SH**, L. Wang)
   Axion Couplings in Heterotic String Theory '24 (P. Agrawal, M. Nee, M Reig)

4. Non-invertible Peccei-Quinn Symmetry and the Massless Quark Solution to Strong CP Problem '24 (C. Cordova, **SH**, S. Koren)
   Spontaneously Broken (-1)-Form U(1) Symmetry '24 (D. Aloni, E. Garcia-Valdecasas, M. Reece, M. Suzuki)
   High-Quality Axions from Higher-Form Symmetries in Extra Dimensions '24 (N. Craig, M. Kongsore)

5. Nonperturbative effects in the Standard Model with gauged 1-form symmetry '21 (M. Anber, E. Popptiz)
   Fractional-charge hadrons and leptons to tell the Standard Model group apart '24 (R. Alonso, D. Dimakou, M. West)
   The Standard Model Gauge Group, SMEFT, and Generalized Symmetries '24 (H-L. Li, L-X. Xu)

6. A New Solution to the Callan-Rubakov Effect '23 (T. D. Brennan)
   Monopoles, Scattering, and Generalized Symmetries '23 (M. Beest, P. B. Smith, D. Delmastro, Z. Komargodski, D. Tong)
   Fermion-Monopole Scattering in the Standard Model '23 (M. Beest, P. B. Smith, D. Delmastro, R. Mouland, D. Tong)

Outline

I. Generalized Global Symmetries

I-1. Higher-form symmetry
I-2. Non-invertible symmetry

II. Strong CP Problem-I: IR to UV

II-1. Non-invertible Peccei-Quinn symmetry
II-2. Massless quark solution

III. Strong CP Problem-II: UV to IR

III-1. $SU(9)$ Color-Flavor unification
III-2. Flavor structure and CKM CPV phase
III-3. Quality Problem

Outline

I. Generalized Global Symmetries

I-1. Higher-form symmetry

I-2. Non-invertible symmetry

II. Strong CP Problem-I: IR to UV

II-1. Non-invertible Peccei-Quinn symmetry

II-2. Massless quark solution

III. Strong CP Problem-II: UV to IR

III-1. $SU(9)$ Color-Flavor unification

III-2. Flavor structure and CKM CPV phase


III-3. Quality Problem

Higher-form symmetries


Various **extended objects** appear in broad class of theories.




Local operator
e.g. particle
0-form symmetry



Line operator
e.g. Wilson line
't Hooft line
1-form symmetry



Surface operator
e.g. Cosmic string
2-form symmetry



Volume operator
e.g. Domain Wall
3-form symmetry

Higher-form symmetries

1. p-form symmetry

0-form $\leftrightarrow$ local op (particle)
0-form $\leftrightarrow$ j_1 (j_μ)
0-form $\leftrightarrow$ A_1 (A_μ)

$$S \supset \int d^4x\, A_\mu j^\mu = \int A_1 \wedge * j_1$$

$$U(\alpha, \Sigma_3) = e^{i\alpha \int * j_1}$$

p-form $\leftrightarrow$ p-dim op
p-form $\leftrightarrow$ j_{p+1}
p-form $\leftrightarrow$ A_{p+1}

$$S \supset \int A_{p+1} \wedge * j_{p+1}$$

$$U\left(\alpha, \Sigma_{d-p-1}\right) = e^{i\alpha \int * j_{p+1}}$$

Higher-form symmetries

1. p-form symmetry

0-form $\leftrightarrow$ local op (particle)
0-form $\leftrightarrow$ j_1 (j_μ)
0-form $\leftrightarrow$ A_1 (A_μ)

p-form $\leftrightarrow$ p-dim op
p-form $\leftrightarrow$ j_{p+1}
p-form $\leftrightarrow$ A_{p+1}

$$S \supset \int d^4x\, A_\mu j^\mu = \int A_1 \wedge * j_1$$

$$S \supset \int A_{p+1} \wedge * j_{p+1}$$

$$U(\alpha, \Sigma_3) = e^{i\alpha \int * j_1}$$


$$U\left(\alpha, \Sigma_{d-p-1}\right) = e^{i\alpha \int * j_{p+1}}$$

"Symmetry Defect Operator"

$$Q(\Sigma_3) = \int_{\Sigma_3} d^3x\, J^0 = \int_{\Sigma_3} * J_1$$

$$U(\alpha, \Sigma_3) = e^{i\alpha Q(\Sigma_3)}$$

$$\langle U(\alpha, \Sigma_3)\psi(x)\rangle \sim e^{i\alpha}\psi(x)$$



Higher-form symmetries

1. p-form symmetry

0-form $\leftrightarrow$ local op (particle)
0-form $\leftrightarrow$ j_1 (j_μ)
0-form $\leftrightarrow$ A_1 (A_μ)

p-form $\leftrightarrow$ p-dim op
p-form $\leftrightarrow$ j_{p+1}
p-form $\leftrightarrow$ A_{p+1}

$$S \supset \int d^4x\, A_\mu j^\mu = \int A_1 \wedge * j_1$$

$$S \supset \int A_{p+1} \wedge * j_{p+1}$$

$$U(\alpha, \Sigma_3) = e^{i\alpha \int * j_1}$$


$$U\left(\alpha, \Sigma_{d-p-1}\right) = e^{i\alpha \int * j_{p+1}}$$

"Symmetry Defect Operator"

$$Q(\Sigma_3) = \int_{\Sigma_3} d^3x\, J^0 = \int_{\Sigma_3} * J_1$$

$$U(\alpha, \Sigma_3) = e^{i\alpha Q(\Sigma_3)}$$

$$\langle U(\alpha, \Sigma_3)\psi(x)\rangle \sim e^{i\alpha}\psi(x)$$



$$\Delta Q = \int_{\tilde{\Sigma}_4} d * J_1 = 0$$

$U(\alpha, \Sigma_3)$ = topological

Higher-form symmetries

1. p-form symmetry

0-form $\leftrightarrow$ local op (particle)
0-form $\leftrightarrow$ j_1 (j_μ)
0-form $\leftrightarrow$ A_1 (A_μ)

p-form $\leftrightarrow$ p-dim op
p-form $\leftrightarrow$ j_{p+1}
p-form $\leftrightarrow$ A_{p+1}


$$S \supset \int d^4x\, A_\mu j^\mu = \int A_1 \wedge * j_1$$


$$S \supset \int A_{p+1} \wedge * j_{p+1}$$

$$U(\alpha, \Sigma_3) = e^{i\alpha \int * j_1}$$

$$U\left(\alpha, \Sigma_{d-p-1}\right) = e^{i\alpha \int * j_{p+1}}$$

E.g.) 0- and 1-form symmetry in 3d





Higher-form symmetries

1. p-form symmetry

1-1. $SU(N)$ YM (either pure YM or with only adj matter)

$\exists\ Z_N^{(1)}(e)$: under 0-form center $\Psi \to e^{\frac{2\pi i}{N} * N}\ \Psi$

$\to$ Wilson line with charge $= 0,1,\cdots,(N-1)$ not screened

$\nexists$ mag 1-form : $\Pi_1\left(SU(N)\right) = \emptyset$

Higher-form symmetries

1. p-form symmetry

1-1. $SU(N)$ YM (either pure YM or with only adj matter)

$\exists\, Z_N^{(1)}(e)$: under 0-form center $\Psi \to e^{\frac{2\pi i}{N} * N}\, \Psi$

$\to$ Wilson line with charge $= 0,1,\cdots,(N-1)$ not screened

$\nexists$ mag 1-form : $\Pi_1\left(SU(N)\right) = \emptyset$

1-2. $PSU(N) = \frac{SU(N)}{Z_N}$: $Z_N^{(1)}(e)$ is gauged (electric states projected out)

$\nexists$ electric 1-form

$\exists\, Z_N^{(1)}(m) : \Pi_1\left(PSU(N)\right) = Z_N$ or $"N * \frac{1}{N} = 1"$

$\Rightarrow \oint G_2 = 2\pi/N\,, \quad \int \mathrm{tr}(G_2 \wedge G_2) \supset \frac{N-1}{2N} \int w_2 \wedge w_2$

Fractional Instanton

Outline

I. Generalized Global Symmetries

I-1. Higher-form symmetry
I-2. Non-invertible symmetry

II. Strong CP Problem-I: IR to UV

II-1. Non-invertible Peccei-Quinn symmetry
II-2. Massless quark solution

III. Strong CP Problem-II: UV to IR

III-1. $SU(9)$ Color-Flavor unification
III-2. Flavor structure and CKM CPV phase
III-3. Quality Problem

Non-Invertible Symmetry

From Fractional Instanton

e.g. $G = SU(N)$

electric 1-form: Z_N

magnetic 1-form: none

Non-Invertible Symmetry

From Fractional Instanton

e.g. $G = SU(N)/Z_L$

electric 1-form: $Z_{N/L}$

magnetic 1-form: Z_L

Non-Invertible Symmetry

From Fractional Instanton

e.g. $G = SU(N)/Z_L$ electric 1-form: $Z_{N/L}$

magnetic 1-form: Z_L

$U(1)_A$ with $\alpha = \frac{2\pi}{z}$, $S \to S + \frac{2\pi K i}{z}\underbrace{\int_{M_4} \frac{G\wedge G}{8\pi^2}}_{\in Z} + \frac{2\pi K i}{z}\underbrace{\left(\frac{L-1}{L}\right)\int_{M_4} \frac{w_2 \wedge w_2}{2}}_{\in Z_L}$

Non-Invertible Symmetry


From Fractional Instanton

e.g. $G = SU(N)/Z_L$ electric 1-form: $Z_{N/L}$

magnetic 1-form: Z_L

$U(1)_A$ with $\alpha = \frac{2\pi}{z}$, $S \to S + \frac{2\pi K i}{z}\int_{M_4}\frac{G\wedge G}{8\pi^2} + \frac{2\pi K i}{z}\left(\frac{L-1}{L}\right)\int_{M_4}\frac{w_2\wedge w_2}{2}$

$\in Z$ $\in Z_L$



Non-Invertible Symmetry


From Fractional Instanton

e.g. $G = SU(N)/Z_L$ electric 1-form: $Z_{N/L}$

magnetic 1-form: Z_L

$U(1)_A$ with $\alpha = \frac{2\pi}{z}$, $S \to S + \frac{2\pi K i}{z}\underbrace{\int_{M_4} \frac{G\wedge G}{8\pi^2}}_{\in Z} + \frac{2\pi K i}{z}\underbrace{\left(\frac{L-1}{L}\right)\int_{M_4} \frac{w_2 \wedge w_2}{2}}_{\in Z_L}$

Global $U(1)_A$



→ **Z Instanton**

Non-Invertible Symmetry


From Fractional Instanton

e.g. $G = SU(N)/Z_L$ electric 1-form: $Z_{N/L}$

magnetic 1-form: Z_L

$U(1)_A$ with $\alpha = \frac{2\pi}{z}$, $S \to S + \frac{2\pi K i}{z}\underbrace{\int_{M_4} \frac{G\wedge G}{8\pi^2}}_{\in Z} + \frac{2\pi K i}{z}\underbrace{\left(\frac{L-1}{L}\right)\int_{M_4} \frac{w_2 \wedge w_2}{2}}_{\in Z_L}$

Global $U(1)_A$



→ **Z Instanton**

→ **Z_L (fractional) Instanton**


Non-Invertible Symmetry

From Fractional Instanton

e.g. $G = SU(N)/Z_L$ electric 1-form: $Z_{N/L}$

magnetic 1-form: Z_L

$U(1)_A$ with $\alpha = \frac{2\pi}{z}$, $S \to S + \frac{2\pi K i}{z}\underbrace{\int_{M_4} \frac{G\wedge G}{8\pi^2}}_{\in Z} + \frac{2\pi K i}{z}\underbrace{\left(\frac{L-1}{L}\right)\int_{M_4} \frac{w_2 \wedge w_2}{2}}_{\in Z_L}$




→ Z Instanton

→ Z_L (fractional) Instanton

Non-Invertible Symmetry

From Fractional Instanton

$$U(1)^{(0)}_A : \quad \Psi_+ \to e^{i\alpha}\,\Psi_+ \,, \quad \Psi_- \to e^{i\alpha}\,\Psi_- \quad \Rightarrow \quad d * j_A \propto w_2 \wedge w_2$$




$$S \;\to\; S + \frac{2\pi i K\ell}{K}\left(\frac{L-1}{L}\right)\int \frac{w_2 \wedge w_2}{2}$$

$$\underbrace{\exp\left(\frac{2\pi\ell i}{K}\oint * j_1\right)}_{U\left(\frac{2\pi\ell}{K}, \Sigma_3\right)}$$

Non-Invertible Symmetry

From Fractional Instanton

$$U(1)^{(0)}_A : \quad \Psi_+ \to e^{i\alpha}\,\Psi_+ \,, \quad \Psi_- \to e^{i\alpha}\,\Psi_- \quad \Rightarrow \quad d * j_A \propto w_2 \wedge w_2$$



$$S \;\to\; S + \frac{2\pi i K\ell}{K}\left(\frac{L-1}{L}\right)\int \frac{w_2 \wedge w_2}{2} - \frac{2\pi i p}{L}\int \frac{w_2 \wedge w_2}{2} \;\to\; S$$


$$\underbrace{\exp\left(\frac{2\pi\ell i}{K}\oint * j_1\right)}_{U\left(\frac{2\pi\ell}{K}, \Sigma_3\right)} \times \mathcal{A}^{\mathrm{L,p}}(\mathrm{w}_2)$$

$$S_{3d} = \frac{iL}{4\pi}\int_{\Sigma_3} a_1 \wedge da_2 + \frac{i}{2\pi}\int_{\Sigma_3} a_1 \wedge w_2 \quad (\text{for } p = 1)$$

Non-Invertible Symmetry

From Fractional Instanton

$$U(1)^{(0)}_A : \quad \Psi_+ \to e^{i\alpha}\,\Psi_+ \ , \quad \Psi_- \to e^{i\alpha}\,\Psi_- \quad \Rightarrow \quad d * j_A \propto w_2 \wedge w_2$$



$$S \ \to \ S + \frac{2\pi i K \ell}{K}\left(\frac{L-1}{L}\right)\int \frac{w_2 \wedge w_2}{2} - \frac{2\pi i p}{L}\int \frac{w_2 \wedge w_2}{2} \ \to \ S$$

$$\boldsymbol{\mathcal{D}_\ell(\Sigma_3)} = \underbrace{\exp\left(\frac{2\pi \ell i}{K}\oint * j_1\right)}_{U\left(\frac{2\pi\ell}{K}, \Sigma_3\right)} \times \mathcal{A}^{\mathrm{L,p}}(\mathrm{w_2})$$

$$\mathcal{D}_\ell(\Sigma_3) \times \overline{\mathcal{D}}_\ell(\Sigma_3) \sim \sum_S \xi(S) \exp\left(\frac{i}{2\pi L}\int_s w_2\right) \neq 1$$

$$S_{3d} = \frac{iL}{4\pi}\int_{\Sigma_3} a_1 \wedge da_2 + \frac{i}{2\pi}\int_{\Sigma_3} a_1 \wedge w_2 \quad (\text{for } p = 1)$$

Outline

I. Generalized Global Symmetries

I-1. Higher-form symmetry
I-2. Non-invertible symmetry

II. Strong CP Problem-I: IR to UV


II-1. Non-invertible Peccei-Quinn symmetry
II-2. Massless quark solution

III. Strong CP Problem-II: UV to IR

III-1. $SU(9)$ Color-Flavor unification
III-2. Flavor structure and CKM CPV phase
III-3. Quality Problem


Strong-CP Problem

1. Strong CP Problem




Strong-CP Problem

1. Strong CP Problem



Strong-CP Problem

1. Strong CP Problem




Expectation based on general rules of effective field theory

$$S_{QCD} \supset \frac{i\bar{\theta}}{8\pi^2}\int Tr(G \wedge G)$$

$$\bar{\theta} \sim O(1)$$

Strong-CP Problem

1. Strong CP Problem



Expectation based on general rules of effective field theory


$$S_{QCD} \supset \frac{i\bar{\theta}}{8\pi^2}\int Tr(G \wedge G)$$

Neutron Electric Dipole Moment

$$d_n \sim 3 \times 10^{16}\,\bar{\theta} \;\rightarrow\; \bar{\theta} < 10^{-10}$$

Strong-CP Problem

1. Strong CP Problem



Expectation based on general rules of effective field theory

$$S_{QCD} \supset \frac{i\bar{\theta}}{8\pi^2}\int Tr(G \wedge G)$$

Neutron Electric Dipole Moment

$$d_n \sim 3 \times 10^{16}\,\bar{\theta} \;\rightarrow\; \bar{\theta} < 10^{-10}$$

$\tilde{J} = \mathbf{Im}\,\mathbf{det}[y_u^\dagger y_u, y_d^\dagger y_d] \propto \sin\delta_{CKM}$ "Jarlskog invariant"

vs

$\bar{\theta} \equiv \arg e^{-i\theta}\,\det(y_u y_d)$

Strong-CP Problem

2. Non-invertible Peccei-Quinn Symmetry

Conclusion:

We start with $\mathcal{L} \supset y_u \widetilde{H} Q \bar{u} + y_e H L \bar{e}$ but $\boldsymbol{y_d = 0}$ $(y_d H Q \bar{d})$

So, new symmetries appearing below are approximate symmetries and $\boldsymbol{y_d}$ is the symmetry breaking spurion (parameter).

Strong-CP Problem

2. Non-invertible Peccei-Quinn Symmetry


Conclusion:

We start with $\mathcal{L} \supset y_u \tilde{H} Q \bar{u} + y_e H L \bar{e}$ but $\boldsymbol{y_d = 0}$ $(y_d H Q \bar{d})$

So, new symmetries appearing below are approximate symmetries and $\boldsymbol{y_d}$ is the symmetry breaking spurion (parameter).

(1) $SU(3)_C \times SU(3)_H / Z_3$: $\boldsymbol{Z_3^{\bar{d}}}$ NIS

(2) $SU(3)_C \times U(1)_H / Z_3,\ H = B_1 + B_2 - 2B_3$: $\boldsymbol{Z_3^{Q-\bar{u}+\bar{d}}}$ NIS



$B_i \equiv Q_i - \bar{u}_i - \bar{d}_i$

Strong-CP Problem

2. Non-invertible Peccei-Quinn Symmetry


Conclusion:

We start with $\mathcal{L} \supset y_u \tilde{H} Q \bar{u} + y_e H L \bar{e}$ but $\boldsymbol{y_d = 0}$ $(y_d H Q \bar{d})$

So, new symmetries appearing below are approximate symmetries and $\boldsymbol{y_d}$ is the symmetry breaking spurion (parameter).

(1) $SU(3)_C \times SU(3)_H / \boldsymbol{Z_3}$: $\boxed{N_c = N_g}$ $\boldsymbol{Z_3^{\bar{d}}}$ NIS

(2) $SU(3)_C \times U(1)_H / Z_3,\ H = B_1 + B_2 - 2B_3$: $\boldsymbol{Z_3^{Q-\bar{u}+\bar{d}}}$ NIS



$B_i \equiv Q_i - \bar{u}_i - \bar{d}_i$

Strong-CP Problem

2. Non-invertible Peccei-Quinn Symmetry

Conclusion:


We start with $\mathcal{L} \supset y_u \tilde{H} Q \bar{u} + y_e H L \bar{e}$ but $\boldsymbol{y_d = 0}$ $(y_d H Q \bar{d})$

So, new symmetries appearing below are approximate symmetries

and $\boldsymbol{y_d}$ is the symmetry breaking spurion (parameter).

(1) $SU(3)_C \times SU(3)_H / Z_3$: $\boldsymbol{Z_3^{\bar{d}}}$ NIS

(2) $SU(3)_C \times U(1)_H / \boldsymbol{Z_3}$, $H = B_1 + B_2 - 2B_3$: $\boldsymbol{Z_3^{Q-\bar{u}+\bar{d}}}$ NIS



$[Z_3^C \subset SU(3)_C] = [Z_3^H \subset U(1)_H]$: $N_C = N_g$

Strong-CP Problem

2. Non-invertible Peccei-Quinn Symmetry


Conclusion:

We start with $\mathcal{L} \supset y_u \tilde{H} Q \bar{u} + y_e H L \bar{e}$ but $\boldsymbol{y_d = 0}$ $(y_d H Q \bar{d})$

So, new symmetries appearing below are approximate symmetries and $\boldsymbol{y_d}$ is the symmetry breaking spurion (parameter).


(1) $SU(3)_C \times SU(3)_H / Z_3$: $\boldsymbol{Z_3^{\bar{d}}}$ NIS

(2) $SU(3)_C \times U(1)_H / Z_3$, $H = B_1 + B_2 - 2B_3$: $Z_3^{Q-\bar{u}+\bar{d}}$ NIS




$B_i \equiv Q_i - \bar{u}_i - \bar{d}_i$

Solving Strong CP with Non-Invertible Symmetry



- Start with only $y_u \tilde{H} Q \bar{u} \quad (y_d = 0)$
- $Z_3^{\bar{d}}$ **NIS** from fractional CFU instantons
- $y_d H Q \bar{d}$ protected by **NIS**
- $\bar{\theta} =$ unphysical

Solving Strong CP with Non-Invertible Symmetry



$SU(9)$

$\langle\Phi\rangle$

$SU(3)_C \times SU(3)_H/Z_3$


$\langle\Sigma\rangle$

$\mathcal{L}_{UV} \supset y_u \tilde{H} Q \bar{u} + y_u^* e^{-\frac{2\pi}{\alpha_9}} H Q \bar{d} + \cdots$

with $\bar{\theta} = 0$

- Start with only $y_u \tilde{H} Q \bar{u} \quad (y_d = 0)$
- $Z_3^{\bar{d}}$ **NIS** from fractional CFU instantons
- $y_d H Q \bar{d}$ protected by **NIS**
- $\bar{\theta} =$ unphysical

Solving Strong CP with Non-Invertible Symmetry



$SU(9)$

$\langle\Phi\rangle$

$SU(3)_C \times SU(3)_H/Z_3$

$\langle\Sigma\rangle$

SM

$\mathcal{L}_{UV} \supset y_u \tilde{H} Q \bar{u} + y_u^* e^{-\frac{2\pi}{\alpha_9}} H Q \bar{d} + \cdots$

with $\bar{\theta} = 0$

- Start with only $y_u \tilde{H} Q \bar{u}$ $(y_d = 0)$
- $Z_3^{\bar{d}}$ **NIS** from fractional CFU instantons
- $y_d H Q \bar{d}$ protected by **NIS**
- $\bar{\theta} =$ unphysical

Yukawa texture and CKM CPV (+possible $\bar{\theta}$)

with $\bar{\theta} = 0$ or $\bar{\theta} \ll 1$

Strong-CP Problem

2. Non-invertible Peccei-Quinn Symmetry

(1) $SU(3)_C \times SU(3)_H/Z_3$: $\boldsymbol{Z_3^{\bar{d}}}$ NIS

	$SU(3)_C$	$SU(3)_H$	$U(1)_B$	$U(1)_{\bar{d}}$
$\boldsymbol{Q}$	3	3	$+1$	0
$\boldsymbol{\bar{u}}$	$\bar{3}$	$\bar{3}$	-1	0
$\boldsymbol{\bar{d}}$	$\bar{3}$	$\bar{3}$	-1	$+1$

$\mathcal{L} \sim y_u \tilde{H} Q^i \bar{u}_i$ (flavor-diagonal/universal)

$y_u = c_u \mathbb{I}_3$, $\mathrm{c_u} =$ number

Strong-CP Problem

2. Non-invertible Peccei-Quinn Symmetry

(1) $SU(3)_C \times SU(3)_H/Z_3$: $\boldsymbol{Z_3^{\bar{d}}}$ NIS

	$SU(3)_C$	$SU(3)_H$	$U(1)_B$	$U(1)_{\bar{d}}$
$\boldsymbol{Q}$	3	3	$+1$	0
$\bar{\boldsymbol{u}}$	$\bar{3}$	$\bar{3}$	-1	0
$\bar{\boldsymbol{d}}$	$\bar{3}$	$\bar{3}$	-1	$+1$

$\mathcal{L} \sim y_u \tilde{H} Q^i \bar{u}_i$ (flavor-diagonal/universal)

$y_u = c_u \mathbb{I}_3$, $\mathrm{c_u} =$ number

CFU Fractional Instanton (**CFU**=**C**olor-(non-abelian)**F**lavor-**U**(1))

Quotient by Z_3: (i) $[Z_3 \in SU(3)_C] \equiv [Z_3 \in SU(3)_H]$

(ii) Under "diagonal" Z_3 entire fields are neutral, more magnetic states

(iii) $\exists Z_3$ magnetic 1-form: $\oint w_2(C) = \oint w_2(H) = 0,1,2 \ (\in Z_3)$

(iv) CFU instanton: $\mathcal{N}_C = \frac{1}{3}\int w_2(C) \wedge w_2(C)$, $\mathcal{N}_H = \frac{1}{3}\int w_2(H) \wedge w_2(H)$

Strong-CP Problem

2. Non-invertible Peccei-Quinn Symmetry

(1) $SU(3)_C \times SU(3)_H/Z_3$: $\boldsymbol{Z_3^{\bar{d}}}$ NIS


	$SU(3)_C$	$SU(3)_H$	$U(1)_B$	$U(1)_{\bar{d}}$
$\boldsymbol{Q}$	3	3	$+1$	0
$\boldsymbol{\bar{u}}$	$\bar{3}$	$\bar{3}$	-1	0
$\boldsymbol{\bar{d}}$	$\bar{3}$	$\bar{3}$	-1	$+1$

$\mathcal{L} \sim y_u \tilde{H} Q^i \bar{u}_i$ (flavor-diagonal/universal)

$y_u = c_u \mathbb{I}_3$, $\mathrm{c_u} =$ number

CFU Fractional Instanton (**CFU**=**C**olor-(non-abelian)**F**lavor-**U**(1))

Quotient by Z_3: (i) $[Z_3 \in SU(3)_C] \equiv [Z_3 \in SU(3)_H]$



(ii) Under "diagonal" Z_3 entire fields are neutral, more magnetic states

(iii) $\exists Z_3$ magnetic 1-form: $\oint w_2(C) = \oint w_2(H) = 0,1,2 \ (\in Z_3)$

(iv) CFU instanton: $\mathcal{N}_C = \frac{1}{3}\int w_2(C) \wedge w_2(C)$, $\mathcal{N}_H = \frac{1}{3}\int w_2(H) \wedge w_2(H)$

$$\mathcal{A}_f = \sum_{\Psi_i} q_i I_{\Psi_i} = 3(\mathcal{N}_C + \mathcal{N}_H)(2q_Q + q_{\bar{d}} + q_{\bar{u}})$$

Strong-CP Problem

2. Non-invertible Peccei-Quinn Symmetry

(1) $SU(3)_C \times SU(3)_H/Z_3$: $\boldsymbol{Z_3^{\bar{d}}}$ NIS

	$SU(3)_C$	$SU(3)_H$	$U(1)_B$	$U(1)_{\bar{d}}$
$\boldsymbol{Q}$	3	3	$+1$	0
$\bar{\boldsymbol{u}}$	$\bar{3}$	$\bar{3}$	-1	0
$\bar{\boldsymbol{d}}$	$\bar{3}$	$\bar{3}$	-1	$+1$

$\mathcal{L} \sim y_u \tilde{H} Q^i \bar{u}_i$ (flavor-diagonal/universal)

$y_u = c_u \mathbb{I}_3$, $\mathrm{c_u} =$ number

	$U(1)_B$	$U(1)_{\bar{d}}$
$[SU(3)_C]^2$	0	N_g
$[SU(2)_L]^2$	$N_c N_g$	0
$[U(1)_Y]^2$	$-18 N_c N_g$	$4 N_c N_g$
$[SU(3)_H]^2$	0	N_c
$[CH]$	0	2

Strong-CP Problem

2. Non-invertible Peccei-Quinn Symmetry

(1) $SU(3)_C \times SU(3)_H/Z_3$: $\boldsymbol{Z_3^{\bar{d}}}$ NIS

	$SU(3)_C$	$SU(3)_H$	$U(1)_B$	$U(1)_{\bar{d}}$
$\boldsymbol{Q}$	3	3	$+1$	0
$\boldsymbol{\bar{u}}$	$\bar{3}$	$\bar{3}$	-1	0
$\boldsymbol{\bar{d}}$	$\bar{3}$	$\bar{3}$	-1	$+1$

$\mathcal{L} \sim y_u \tilde{H} Q^i \bar{u}_i$ (flavor-diagonal/universal)

$y_u = c_u \mathbb{I}_3$, $\mathrm{c_u} =$ number

	$U(1)_B$	$U(1)_{\bar{d}}$
$[SU(3)_C]^2$	0	N_g
$[SU(2)_L]^2$	$N_c N_g$	0
$[U(1)_Y]^2$	$-18 N_c N_g$	$4 N_c N_g$
$[SU(3)_H]^2$	0	N_c
$[CH]$	0	2

Symmetry

(i) Without including $[CH]$ instanton:

$\frac{U(1)_B}{Z_3} \times U(1)_{\bar{d}} \to Z_3^B \times Z_3^{\bar{d}}$

non-abelian instantons dominant → No NIS

Strong-CP Problem

2. Non-invertible Peccei-Quinn Symmetry

(1) $SU(3)_C \times SU(3)_H/Z_3$: $\boldsymbol{Z_3^{\bar{d}}}$ NIS

	$SU(3)_C$	$SU(3)_H$	$U(1)_B$	$U(1)_{\bar{d}}$
$\boldsymbol{Q}$	3	3	$+1$	0
$\bar{\boldsymbol{u}}$	$\bar{3}$	$\bar{3}$	-1	0
$\bar{\boldsymbol{d}}$	$\bar{3}$	$\bar{3}$	-1	$+1$

$\mathcal{L} \sim y_u \tilde{H} Q^i \bar{u}_i$ (flavor-diagonal/universal)

$y_u = c_u \mathbb{I}_3, \quad c_u =$ number

	$U(1)_B$	$U(1)_{\bar{d}}$
$[SU(3)_C]^2$	0	N_g
$[SU(2)_L]^2$	$N_c N_g$	0
$[U(1)_Y]^2$	$-18 N_c N_g$	$4 N_c N_g$
$[SU(3)_H]^2$	0	N_c
$[CH]$	0	2

Symmetry

(i) Without including $[CH]$ instanton:

$\frac{U(1)_B}{Z_3} \times U(1)_{\bar{d}} \to Z_3^B \times Z_3^{\bar{d}}$

non-abelian instantons dominant → No NIS

(ii) With $[CH]$ instanton:

$Z_3^B \times Z_3^{\bar{d}} \to Z_3^B \times \emptyset$

Strong-CP Problem

2. Non-invertible Peccei-Quinn Symmetry

(1) $SU(3)_C \times SU(3)_H/Z_3$: $\boldsymbol{Z_3^{\bar{d}}}$ NIS

	$SU(3)_C$	$SU(3)_H$	$U(1)_B$	$U(1)_{\bar{d}}$
$\boldsymbol{Q}$	3	3	$+1$	0
$\bar{\boldsymbol{u}}$	$\bar{3}$	$\bar{3}$	-1	0
$\bar{\boldsymbol{d}}$	$\bar{3}$	$\bar{3}$	-1	$+1$

$\mathcal{L} \sim y_u \tilde{H} Q^i \bar{u}_i$ (flavor-diagonal/universal)

$y_u = c_u \mathbb{I}_3, \quad c_u =$ number

	$U(1)_B$	$U(1)_{\bar{d}}$
$[SU(3)_C]^2$	0	N_g
$[SU(2)_L]^2$	$N_c N_g$	0
$[U(1)_Y]^2$	$-18 N_c N_g$	$4 N_c N_g$
$[SU(3)_H]^2$	0	N_c
$[CH]$	0	2

Symmetry

(i) Without including $[CH]$ instanton:

$\frac{U(1)_B}{Z_3} \times U(1)_{\bar{d}} \to Z_3^B \times Z_3^{\bar{d}}$

non-abelian instantons dominant → No NIS

(ii) With $[CH]$ instanton:

$Z_3^B \times Z_3^{\bar{d}} \to Z_3^B \times \emptyset$

Strong-CP Problem


2. Non-invertible Peccei-Quinn Symmetry

(1) $SU(3)_C \times SU(3)_H/Z_3$: $\boldsymbol{Z_3^{\bar{d}}}$ NIS

	$SU(3)_C$	$SU(3)_H$	$U(1)_B$	$U(1)_{\bar{d}}$
$\boldsymbol{Q}$	3	3	+1	0
$\boldsymbol{\bar{u}}$	$\bar{3}$	$\bar{3}$	−1	0
$\boldsymbol{\bar{d}}$	$\bar{3}$	$\bar{3}$	−1	+1

$\mathcal{L} \sim y_u \tilde{H} Q^i \bar{u}_i$ (flavor-diagonal/universal)

$y_u = c_u \mathbb{I}_3$, $c_u =$ number



Symmetry

(i) Without including $[CH]$ instanton:

$\frac{U(1)_B}{Z_3} \times U(1)_{\bar{d}} \rightarrow Z_3^B \times Z_3^{\bar{d}}$

non-abelian instantons dominant → No NIS

(ii) With $[CH]$ instanton:

$Z_3^B \times Z_3^{\bar{d}} \rightarrow Z_3^B \times \emptyset \quad \Rightarrow \quad \boldsymbol{Z_3^{\bar{d}}}$ **NIPQ Symmetry**

Strong-CP Problem


2. Non-invertible Peccei-Quinn Symmetry

(1) $SU(3)_C \times SU(3)_H/Z_3$: $\boldsymbol{Z_3^{\bar{d}}}$ NIS

	$SU(3)_C$	$SU(3)_H$	$U(1)_B$	$U(1)_{\bar{d}}$
$\boldsymbol{Q}$	3	3	+1	0
$\boldsymbol{\bar{u}}$	$\bar{3}$	$\bar{3}$	−1	0
$\boldsymbol{\bar{d}}$	$\bar{3}$	$\bar{3}$	−1	+1

$\mathcal{L} \sim y_u \tilde{H} Q^i \bar{u}_i$ (flavor-diagonal/universal)

$y_u = c_u \mathbb{I}_3$, $c_u =$ number



Symmetry

(i) Without including $[CH]$ instanton:

$$\frac{U(1)_B}{Z_3} \times U(1)_{\bar{d}} \rightarrow Z_3^B \times Z_3^{\bar{d}}$$

non-abelian instantons dominant → No NIS

(ii) With $[CH]$ instanton: $N_c = N_g$

$Z_3^B \times Z_3^{\bar{d}} \rightarrow Z_3^B \times \emptyset \Rightarrow \boldsymbol{Z_3^{\bar{d}}}$ **NIPQ Symmetry**

Outline

I. Generalized Global Symmetries

I-1. Higher-form symmetry
I-2. Non-invertible symmetry

II. Strong CP Problem-I: IR to UV

II-1. Non-invertible Peccei-Quinn symmetry
II-2. Massless quark solution

III. Strong CP Problem-II: UV to IR

III-1. $SU(9)$ Color-Flavor unification
III-2. Flavor structure and CKM CPV phase
III-3. Quality Problem

Strong-CP Problem

3. Massless Quark Solution to the Strong CP Problem

(1) $SU(3)_C \times SU(3)_H/Z_3$: $\boldsymbol{Z_3^{\bar{d}}}$ NIS

$\mathcal{L} \sim y_d H Q \bar{d}$ term is forbidden by $Z_3^{\bar{d}}$ non-invertible Peccei-Quinn symmetry

Down quarks (d, s, b) are massless if $Z_3^{\bar{d}}$ is exact.

(y_d = non-invertible $Z_3^{\bar{d}}$ breaking spurion (paramter))

Strong-CP Problem

3. Massless Quark Solution to the Strong CP Problem

(1) $SU(3)_C \times SU(3)_H/Z_3$: $\boldsymbol{Z_3^{\bar{d}}}$ NIS

$\mathcal{L} \sim y_d H Q \bar{d}$ term is forbidden by $Z_3^{\bar{d}}$ non-invertible Peccei-Quinn symmetry

Down quarks (d, s, b) are massless if $Z_3^{\bar{d}}$ is exact.

(y_d = non-invertible $Z_3^{\bar{d}}$ breaking spurion (paramter))

Massless Quark Solution:

1. $\mathcal{L}_{SM} \supset y_u \tilde{H} Q \bar{u} + y_d H Q \bar{d} + \frac{\theta}{32\pi^2} G\tilde{G}$

$= m_u e^{i\varphi_u} u\bar{u} + m_d e^{i\varphi_d} d\bar{d} + \frac{\theta}{32\pi^2} G\tilde{G}$

$u \to e^{i\alpha} u, \quad \varphi_u \to \varphi_u + \alpha, \quad \theta \to \theta + \alpha$

$d \to e^{i\beta} d, \quad \varphi_d \to \varphi_d + \beta, \quad \theta \to \theta + \beta$

Strong-CP Problem

3. Massless Quark Solution to the Strong CP Problem

(1) $SU(3)_C \times SU(3)_H/Z_3$: $\boldsymbol{Z_3^{\bar{d}}}$ NIS

$\mathcal{L} \sim y_d H Q \bar{d}$ term is forbidden by $Z_3^{\bar{d}}$ non-invertible Peccei-Quinn symmetry

Down quarks (d, s, b) are massless if $Z_3^{\bar{d}}$ is exact.

(y_d = non-invertible $Z_3^{\bar{d}}$ breaking spurion (paramter))

Massless Quark Solution:

1. $\mathcal{L}_{SM} \supset y_u \tilde{H} Q \bar{u} + y_d H Q \bar{d} + \frac{\theta}{32\pi^2} G\tilde{G}$

$= m_u e^{i\varphi_u} u\bar{u} + m_d e^{i\varphi_d} d\bar{d} + \frac{\theta}{32\pi^2} G\tilde{G}$

$u \to e^{i\alpha} u, \quad \varphi_u \to \varphi_u + \alpha, \quad \theta \to \theta + \alpha$

$d \to e^{i\beta} d, \quad \varphi_d \to \varphi_d + \beta, \quad \theta \to \theta + \beta$

$$\boldsymbol{\bar{\theta} \equiv \arg e^{-i\theta} \det(y_u y_d)}$$

Neutron electric dipole moment $d_n \sim 3 \times 10^{16}\, \bar{\theta} \;\to\; \bar{\theta} < 10^{-10}$

Strong-CP Problem

3. Massless Quark Solution to the Strong CP Problem

(1) $SU(3)_C \times SU(3)_H/Z_3$: $\boldsymbol{Z_3^{\bar{d}}}$ NIS

$\mathcal{L} \sim y_d H Q \bar{d}$ term is forbidden by $Z_3^{\bar{d}}$ non-invertible Peccei-Quinn symmetry

Down quarks (d, s, b) are massless if $Z_3^{\bar{d}}$ is exact.

(y_d = non-invertible $Z_3^{\bar{d}}$ breaking spurion (paramter))

Massless Quark Solution:

2. In the presence of massless chiral fermion, e.g. $m_d = 0$, "$\boldsymbol{\bar{\theta}}$ is unphysical"

$$\mathcal{L}_{SM} \supset y_u \tilde{H} Q \bar{u} + \frac{\theta}{32\pi^2} G\tilde{G} = m_u e^{i\varphi_u} u\bar{u} + \frac{\theta}{32\pi^2} G\tilde{G}$$

Field redefinition: $d \to e^{i\beta} d \quad \Rightarrow \quad \delta S = \frac{i}{8\pi^2}(\bar{\theta} + \beta) \int \mathrm{tr}(G_2 \wedge G_2)$

Strong-CP Problem

3. Massless Quark Solution to the Strong CP Problem

(1) $SU(3)_C \times SU(3)_H/Z_3$: $\boldsymbol{Z_3^{\bar{d}}}$ NIS

$\mathcal{L} \sim y_d H Q \bar{d}$ term is forbidden by $Z_3^{\bar{d}}$ non-invertible Peccei-Quinn symmetry

Down quarks (d, s, b) are massless if $Z_3^{\bar{d}}$ is exact.

(y_d = non-invertible $Z_3^{\bar{d}}$ breaking spurion (paramter))

Massless Quark Solution:

2. In the presence of massless chiral fermion, e.g. $m_d = 0$, "$\boldsymbol{\bar{\theta}}$ is unphysical"

$$\mathcal{L}_{SM} \supset y_u \tilde{H} Q \bar{u} + \frac{\theta}{32\pi^2} G\tilde{G} = m_u e^{i\varphi_u} u\bar{u} + \frac{\theta}{32\pi^2} G\tilde{G}$$

Field redefinition: $d \to e^{i\beta} d \quad \Rightarrow \quad \delta S = \frac{i}{8\pi^2}(\bar{\theta} + \beta) \int \mathrm{tr}(G_2 \wedge G_2)$

Note: $M \equiv e^{-i\theta} \det(y_u y_d) \in \mathbb{C}$ and $CP: \mathrm{Im}(M) \to -\mathrm{Im}(M)$

M behaves smoothly as $|M| \to 0$

CP-invariance $\leftrightarrow$ $M \in \mathbb{R}_+$

Strong-CP Problem

3. Massless Quark Solution to the Strong CP Problem

(1) $SU(3)_C \times SU(3)_H/Z_3$: $\boldsymbol{Z_3^{\bar{d}}}$ NIS

$\mathcal{L} \sim y_d H Q \bar{d}$ term is forbidden by $Z_3^{\bar{d}}$ non-invertible Peccei-Quinn symmetry

Down quarks (d, s, b) are massless if $Z_3^{\bar{d}}$ is exact.

(y_d = non-invertible $Z_3^{\bar{d}}$ breaking spurion (paramter))

Massless Quark Solution:

2. In the presence of massless chiral fermion, e.g. , "$\boldsymbol{\bar{\theta}}$ is unphysical"


$$\mathcal{L}_{SM} \supset y_u \tilde{H} Q \bar{u} + \frac{\theta}{32\pi^2} G\tilde{G} = m_u e^{i\varphi_u} u\bar{u} + \frac{\theta}{32\pi^2} G\tilde{G}$$

Field redefinition: $d \to e^{i\beta} d \quad \Rightarrow \quad \delta S = \frac{i}{8\pi^2}(\bar{\theta} + \beta)$.

Note: $M \equiv e^{-i\theta} \det(y_u y_d) \in \mathbb{C}$ and $CP: \mathrm{Im}(M) \to -$

M behaves smoothly as $|M| \to 0$

CP-invariance $\leftrightarrow$ $M \in \mathbb{R}_+$



Strong-CP Problem

3. Massless Quark Solution to the Strong CP Problem

(1) $SU(3)_C \times SU(3)_H/Z_3$: $\boldsymbol{Z_3^{\bar{d}}}$ NIS

$\mathcal{L} \sim y_d H Q \bar{d}$ term is forbidden by $Z_3^{\bar{d}}$ non-invertible Peccei-Quinn symmetry

Down quarks (d, s, b) are massless if $Z_3^{\bar{d}}$ is exact.

(y_d = non-invertible $Z_3^{\bar{d}}$ breaking spurion (paramter))

Massless Quark Solution:

2. In the presence of massless chiral fermion, e.g. , "$\boldsymbol{\bar{\theta}}$ is unphysical"


$$\mathcal{L}_{SM} \supset y_u \tilde{H} Q \bar{u} + \frac{\theta}{32\pi^2} G\tilde{G} = m_u e^{i\varphi_u} u\bar{u} + \frac{\theta}{32\pi^2} G\tilde{G}$$

Field redefinition: $d \to e^{i\beta} d \quad \Rightarrow \quad \delta S = \frac{i}{8\pi^2}(\bar{\theta} + \beta)$.

Note: $M \equiv e^{-i\theta} \det(y_u y_d) \in \mathbb{C}$ and $CP: \mathrm{Im}(M) \to -$

M behaves smoothly as $|M| \to 0$

CP-invariance $\leftrightarrow$ $M \in \mathbb{R}_+$



Strong-CP Problem

3. Massless Quark Solution to the Strong CP Problem

(1) $SU(3)_C \times SU(3)_H/Z_3$: $\boldsymbol{Z_3^{\bar{d}}}$ NIS

$\mathcal{L} \sim y_d H Q \bar{d}$ term is forbidden by $Z_3^{\bar{d}}$ non-invertible Peccei-Quinn symmetry

Down quarks (d, s, b) are massless if $Z_3^{\bar{d}}$ is exact.

(y_d = non-invertible $Z_3^{\bar{d}}$ breaking spurion (paramter))

Massless Quark Solution:

3. In SM, "massless up quark solution" tried.


In nature, up quark seems to be massive

e.g. Chiral-PT + observed hadron mass : $m_u/m_d \sim 0.6$

QCD instanton calculation not under analytic control

Lattice QCD : QCD instanton **not sufficient** in size

Solving Strong CP with Non-Invertible Symmetry



$SU(9)$

$\langle\Phi\rangle$

$SU(3)_C \times SU(3)_H/Z_3$

$\langle\Sigma\rangle$

SM

$\mathcal{L}_{UV} \supset y_u \tilde{H} Q \bar{u} + y_u^* e^{-\frac{2\pi}{\alpha_9}} H Q \bar{d} + \cdots$


with $\bar{\theta} = 0$

- Start with only $y_u \tilde{H} Q \bar{u}$ $(y_d = 0)$
- $Z_3^{\bar{d}}$ **NIS** from fractional CFU instantons
- $y_d H Q \bar{d}$ protected by **NIS**
- $\bar{\theta} =$ unphysical

Yukawa texture and CKM CPV (+possible $\bar{\theta}$)

with $\bar{\theta} = 0$ or $\bar{\theta} \ll 1$

Solving Strong CP with Non-Invertible Symmetry



Small (G/H) Instanton
[NIS breaking]

$SU(9)$

$\langle\Phi\rangle$

$SU(3)_C \times SU(3)_H/Z_3$

$\langle\Sigma\rangle$

SM

$\mathcal{L}_{UV} \supset y_u \tilde{H} Q \bar{u} + \boldsymbol{y_u^* e^{-\frac{2\pi}{\alpha_9}}} HQ\bar{d} + \cdots$

with $\bar{\theta} = 0$

- Start with only $y_u \tilde{H} Q \bar{u}$ $\quad (y_d = 0)$
- $Z_3^{\bar{d}}$ **NIS** from fractional CFU instantons
- $y_d HQ\bar{d}$ protected by **NIS**
- $\bar{\theta} =$ unphysical

Yukawa texture and CKM CPV (+possible $\bar{\theta}$)

with $\bar{\theta} = 0$ or $\bar{\theta} \ll 1$

Outline

I. Generalized Global Symmetries

I-1. Higher-form symmetry
I-2. Non-invertible symmetry

II. Strong CP Problem-I: IR to UV


II-1. Non-invertible Peccei-Quinn symmetry
II-2. Massless quark solution

III. Strong CP Problem-II: UV to IR

III-1. $SU(9)$ **Color-Flavor unification**
III-2. Flavor structure and CKM CPV phase
III-3. Quality Problem

Color-Flavor Unification


1. $SU(9)$ Unification and the Strong CP phase $\overline{\theta}$



$= 9\ (\square)$ of $SU(9)$

Color-Flavor Unification

1. $SU(9)$ Unification and the Strong CP phase $\overline{\theta}$




$= 9\ (\square)\ \text{of}\ SU(9)$

	$SU(9)$	$U(1)_{Q-\bar{u}}$	$U(1)_{\bar{d}}$
$\boldsymbol{Q = (u, d)^t}$	9	$+1$	0
$\boldsymbol{\bar{u}}$	$\bar{9}$	-1	0
$\boldsymbol{\bar{d}}$	$\bar{9}$	0	$+1$
H	1	0	0

Color-Flavor Unification

1. $SU(9)$ Unification and the Strong CP phase $\bar{\theta}$




$= 9\ (\square)$ of $SU(9)$

	$SU(9)$	$U(1)_{Q-\bar{u}+\bar{d}}$
Φ	165 (3S)	0
$\Sigma_{1,2}$	80 (adj)	0
ρ	9	-1
χ	1	0

	$SU(9)$	$U(1)_{Q-\bar{u}}$	$U(1)_{\bar{d}}$
$\boldsymbol{Q} = (\boldsymbol{u}, \boldsymbol{d})^{\boldsymbol{t}}$	9	$+1$	0
$\bar{\boldsymbol{u}}$	$\bar{9}$	-1	0
$\bar{\boldsymbol{d}}$	$\bar{9}$	0	$+1$
H	1	0	0

Color-Flavor Unification


1. $SU(9)$ Unification and the Strong CP phase $\bar{\theta}$



$= 9\ (\square)\ \text{of}\ SU(9)$


	$SU(9)$	$U(1)_{Q-\bar{u}+\bar{d}}$
Φ	165 (3S)	0
$\Sigma_{1,2}$	80 (adj)	0
ρ	9	-1
χ	1	0

	$SU(9)$	$U(1)_{Q-\bar{u}}$	$U(1)_{\bar{d}}$
$\boldsymbol{Q} = (\boldsymbol{u}, \boldsymbol{d})^{\boldsymbol{t}}$	9	$+1$	0
$\bar{\boldsymbol{u}}$	$\bar{9}$	-1	0
$\bar{\boldsymbol{d}}$	$\bar{9}$	0	$+1$
H	1	0	0



Color-Flavor Unification


1. $SU(9)$ Unification and the Strong CP phase $\bar{\theta}$



$= 9\ (\square)\ \text{of}\ SU(9)$


	$SU(9)$	$U(1)_{Q-\bar{u}+\bar{d}}$
Φ	165 (3S)	0
$\Sigma_{1,2}$	80 (adj)	0
ρ	9	-1
χ	1	0

	$SU(9)$	$U(1)_{Q-\bar{u}}$	$U(1)_{\bar{d}}$
$\boldsymbol{Q} = (\boldsymbol{u}, \boldsymbol{d})^{\boldsymbol{t}}$	9	$+1$	0
$\bar{\boldsymbol{u}}$	$\bar{9}$	-1	0
$\bar{\boldsymbol{d}}$	$\bar{9}$	0	$+1$
H	1	0	0



Color-Flavor Unification


1. $SU(9)$ Unification and the Strong CP phase $\bar{\theta}$



$= 9\ (\square)\ \text{of}\ SU(9)$


	$SU(9)$	$U(1)_{Q-\bar{u}+\bar{d}}$
Φ	165 (3S)	0
$\Sigma_{1,2}$	80 (adj)	0
ρ	9	-1
χ	1	0

	$SU(9)$	$U(1)_{Q-\bar{u}}$	$U(1)_{\bar{d}}$
$\boldsymbol{Q} = (\boldsymbol{u}, \boldsymbol{d})^{\boldsymbol{t}}$	9	$+1$	0
$\bar{\boldsymbol{u}}$	$\bar{9}$	-1	0
$\bar{\boldsymbol{d}}$	$\bar{9}$	0	$+1$
H	1	0	0



Color-Flavor Unification

1. $SU(9)$ Unification and the Strong CP phase $\overline{\theta}$



$= 9\ (\square)\ \text{of}\ SU(9)$

	$SU(9)$	$U(1)_{Q-\bar{u}}$	$U(1)_{\bar{d}}$
$\boldsymbol{Q} = (\boldsymbol{u}, \boldsymbol{d})^{\boldsymbol{t}}$	9	$+1$	0
$\overline{\boldsymbol{u}}$	$\bar{9}$	-1	0
$\overline{\boldsymbol{d}}$	$\bar{9}$	0	$+1$
H	1	0	0


$$\mathcal{L}_0 = y_t \tilde{H} \boldsymbol{Q} \overline{\boldsymbol{u}} + h.c. + \frac{i\theta_9}{32\pi^2} \int F\tilde{F}$$

$$y_t \sim O(1)$$

y_d perturbatively protected by $U(1)_{\bar{d}}$

Color-Flavor Unification

1. $SU(9)$ Unification and the Strong CP phase $\bar{\theta}$



$= 9\ (\square)$ of $SU(9)$

	$SU(9)$	$U(1)_{Q-\bar{u}}$	$U(1)_{\bar{d}}$
$\boldsymbol{Q} = (\boldsymbol{u}, \boldsymbol{d})^{\boldsymbol{t}}$	9	$+1$	0
$\bar{\boldsymbol{u}}$	$\bar{9}$	-1	0
$\bar{\boldsymbol{d}}$	$\bar{9}$	0	$+1$
H	1	0	0

$$\mathcal{L}_0 = y_t \tilde{H} \boldsymbol{Q} \bar{\boldsymbol{u}} + h.c. + \frac{i\theta_9}{32\pi^2} \int F\tilde{F}$$

$y_t \sim O(1)$

y_d perturbatively protected by $U(1)_{\bar{d}}$

$U(1)_{Q-\bar{u}}[SU(9)]^2 = U(1)_{\bar{d}}[SU(9)]^2 = 1$

⇒ [Anomaly Free] $U(1)_{B=Q-\bar{u}-\bar{d}}$
[Anomalous] $U(1)_{Q-\bar{u}+\bar{d}}$ or $U(1)_{\bar{d}}$


Color-Flavor Unification

1. $SU(9)$ Unification and the Strong CP phase $\bar{\theta}$

$SU(3)_H$ →, $SU(3)_C$ ↓

$\bar{u}$	$\bar{c}$	$\bar{t}$
$\bar{u}$	$\bar{c}$	$\bar{t}$
$\bar{u}$	$\bar{c}$	$\bar{t}$

$= 9\ (\square)$ of $SU(9)$



	$SU(9)$	$U(1)_{Q-\bar{u}}$	$U(1)_{\bar{d}}$
$\boldsymbol{Q} = (\boldsymbol{u}, \boldsymbol{d})^{\boldsymbol{t}}$	9	$+1$	0
$\bar{\boldsymbol{u}}$	$\bar{9}$	-1	0
$\bar{\boldsymbol{d}}$	$\bar{9}$	0	$+1$
H	1	0	0

$$\mathcal{L}_0 = y_t \tilde{H} \boldsymbol{Q}\bar{\boldsymbol{u}} + h.c. + \frac{i\theta_9}{32\pi^2}\int F\tilde{F}$$

$$+ y_t^* e^{i\theta_9} e^{-\frac{2\pi}{\alpha_9}} H\boldsymbol{Q}\bar{\boldsymbol{d}}$$

$$\boldsymbol{y_d \sim y_t^* e^{i\theta_9} e^{-\frac{2\pi}{\alpha_9(\Lambda_9)}}}$$

$U(1)_{Q-\bar{u}}[SU(9)]^2 = U(1)_{\bar{d}}[SU(9)]^2 = 1$

⇒ [Anomaly Free] $U(1)_{B=Q-\bar{u}-\bar{d}}$

[Anomalous] $U(1)_{Q-\bar{u}+\bar{d}}$ or $U(1)_{\bar{d}}$


Color-Flavor Unification

1. $SU(9)$ Unification and the Strong CP phase $\bar{\theta}$

$SU(3)_H$ →, $SU(3)_C$ ↓

$\bar{u}$	$\bar{c}$	$\bar{t}$
$\bar{u}$	$\bar{c}$	$\bar{t}$
$\bar{u}$	$\bar{c}$	$\bar{t}$

$= 9\ (\square)$ of $SU(9)$



	$SU(9)$	$U(1)_{Q-\bar{u}}$	$U(1)_{\bar{d}}$
$\boldsymbol{Q=(u,d)^t}$	9	+1	0
$\boldsymbol{\bar{u}}$	$\bar{9}$	−1	0
$\boldsymbol{\bar{d}}$	$\bar{9}$	0	+1
H	1	0	0

$$\mathcal{L}_0 = y_t \tilde{H} \boldsymbol{Q\bar{u}} + h.c. + \frac{i\theta_9}{32\pi^2}\int F\tilde{F}$$

$$+ y_t^* e^{i\theta_9} e^{-\frac{2\pi}{\alpha_9}} H\boldsymbol{Q\bar{d}}$$

$$\boldsymbol{y_d \sim y_t^* e^{i\theta_9} e^{-\frac{2\pi}{\alpha_9(\Lambda_9)}}}$$

$$\bar{\theta} = \arg e^{-i\theta_9} \det(y_u y_d) = -\theta_9 + \arg|y_t|^2\, e^{i\theta_9} = 0 \quad \checkmark$$

Color-Flavor Unification

2. $SU(9) \to SU(3)_C \times SU(3)_H / Z_3$

	$SU(9)$	$U(1)_{Q-\bar{u}+\bar{d}}$
Φ	165 (3S)	0
$\Sigma_{1,2}$	80 (adj)	0
ρ	9	-1
χ	1	0

(i) SSB by $\langle \Phi^{\mathrm{ABC}} \rangle = \Lambda_9 \epsilon^{abc} \epsilon^{ijk}$

(ii) $9\left(\mathrm{Q}, \bar{u}, \bar{d}, \rho\right) \to \ (3,3)$

(iii) Z_3 Quotient: $Q \to g_C \, Q \, g_H^{\dagger}$

(iv) $165 \to (10,10) + (\mathbf{8},\mathbf{8}) + (1,1)$

$(\Phi_{\{ai,bj,ck\}} \sim \Phi_{\{abc\}} \cdot \tilde{\phi}_{\{ijk\}} + \Phi_{[ab]c} \cdot \tilde{\phi}_{[ij]k} + \Phi_{[abc]} \cdot \tilde{\phi}_{[ijk]})$

(v) $80 \to (8,8) + (8,1) + (1,8)$

Color-Flavor Unification

2. $SU(9) \to SU(3)_C \times SU(3)_H/Z_3$

	$SU(9)$	$U(1)_{Q-\bar{u}+\bar{d}}$
Φ	165 (3S)	0
$\Sigma_{1,2}$	80 (adj)	0
ρ	9	-1
χ	1	0

(i) SSB by $\langle \Phi^{\mathrm{ABC}} \rangle = \Lambda_9 \epsilon^{abc} \epsilon^{ijk}$

(ii) $9\left(\mathrm{Q}, \bar{u}, \bar{d}, \rho\right) \to \ (3,3)$

(iii) Z_3 Quotient: $Q \to g_C \, Q \, g_H^\dagger$

(iv) $165 \to (10,10) + (\mathbf{8},\mathbf{8}) + (1,1)$

$(\Phi_{\{ai,bj,ck\}} \sim \Phi_{\{abc\}} \cdot \tilde{\phi}_{\{ijk\}} + \Phi_{[ab]c} \cdot \tilde{\phi}_{[ij]k} + \Phi_{[abc]} \cdot \tilde{\phi}_{[ijk]})$

(v) $80 \to (8,8) + (8,1) + (1,8)$

$$\mathcal{L}_0 = y_t \tilde{H} \boldsymbol{Q}\bar{\boldsymbol{u}} + y_t^* e^{i\theta_9} e^{-\frac{2\pi}{\alpha_9}} H \boldsymbol{Q}\bar{\boldsymbol{d}} + \frac{i\theta_9}{32\pi^2} \int F\tilde{F}$$ (Yukawa=single number)

$$\to y_t \tilde{H} \boldsymbol{Q}\bar{\boldsymbol{u}} + y_t^* e^{i\theta_9} e^{-\frac{2\pi}{3\alpha_S(\Lambda_9)}} H \boldsymbol{Q}\bar{\boldsymbol{d}} + \frac{i3\theta_9}{32\pi^2} \int \left(G\tilde{G} + K\tilde{K}\right)$$ (Yukawa$\propto \mathbb{I}_3$, Flavor-diag)

Index of embedding

$G \to H$: $(1 - H - \text{instanton}) = (n - G - \text{instanton})$

Color-Flavor Unification

2. $SU(9) \to SU(3)_C \times SU(3)_H/Z_3$

	$SU(9)$	$U(1)_{Q-\bar{u}+\bar{d}}$
Φ	165 (3S)	0
$\Sigma_{1,2}$	80 (adj)	0
ρ	9	-1
χ	1	0

(i) SSB by $\langle \Phi^{\mathrm{ABC}} \rangle = \Lambda_9 \epsilon^{abc} \epsilon^{ijk}$

(ii) $9\left(\mathrm{Q}, \bar{u}, \bar{d}, \rho\right) \to \ (3,3)$

(iii) Z_3 Quotient: $Q \to g_C \, Q \, g_H^\dagger$

(iv) $165 \to (10,10) + (\mathbf{8},\mathbf{8}) + (1,1)$

$(\Phi_{\{ai,bj,ck\}} \sim \Phi_{\{abc\}} \cdot \tilde{\phi}_{\{ijk\}} + \Phi_{[ab]c} \cdot \tilde{\phi}_{[ij]k} + \Phi_{[abc]} \cdot \tilde{\phi}_{[ijk]})$

(v) $80 \to (8,8) + (8,1) + (1,8)$

$$\mathcal{L}_0 = y_t \tilde{H} \boldsymbol{Q}\bar{\boldsymbol{u}} + y_t^* e^{i\theta_9} e^{-\frac{2\pi}{\alpha_9}} H \boldsymbol{Q}\bar{\boldsymbol{d}} + \frac{i\theta_9}{32\pi^2} \int F\tilde{F}$$ (Yukawa=single number)

$$\to y_t \tilde{H} \boldsymbol{Q}\bar{\boldsymbol{u}} + y_t^* e^{i\theta_9} e^{-\frac{2\pi}{3\alpha_S(\Lambda_9)}} H \boldsymbol{Q}\bar{\boldsymbol{d}} + \frac{i3\theta_9}{32\pi^2} \int \left(G\tilde{G} + K\tilde{K}\right)$$ (Yukawa$\propto \mathbb{I}_3$, Flavor-diag)

$\bar{\theta} = -3\theta_9 + \arg\det|y_t|^2 \, e^{i\theta_9} = -3\theta_9 + 3\theta_9 = 0$ ✓

Color-Flavor Unification

2. $SU(9) \to SU(3)_C \times SU(3)_H/Z_3$

	$SU(9)$	$U(1)_{Q-\bar{u}+\bar{d}}$
Φ	165 (3S)	0
$\Sigma_{1,2}$	80 (adj)	0
ρ	9	-1
χ	1	0

(i) SSB by $\langle \Phi^{\mathrm{ABC}} \rangle = \Lambda_9 \epsilon^{abc} \epsilon^{ijk}$

(ii) $9\left(\mathrm{Q}, \bar{u}, \bar{d}, \rho\right) \to \ (3,3)$

(iii) Z_3 Quotient: $Q \to g_C \, Q \, g_H^\dagger$

(iv) $165 \to (10,10) + (\mathbf{8},\mathbf{8}) + (1,1)$

$(\Phi_{\{ai,bj,ck\}} \sim \Phi_{\{abc\}} \cdot \tilde{\phi}_{\{ijk\}} + \Phi_{[ab]c} \cdot \tilde{\phi}_{[ij]k} + \Phi_{[abc]} \cdot \tilde{\phi}_{[ijk]})$

(v) $80 \to (8,8) + (8,1) + (1,8)$

$$\mathcal{L}_0 = y_t \tilde{H} \boldsymbol{Q}\bar{\boldsymbol{u}} + y_t^* e^{i\theta_9} e^{-\frac{2\pi}{\alpha_9}} H \boldsymbol{Q}\bar{\boldsymbol{d}} + \frac{i\theta_9}{32\pi^2} \int F\tilde{F}$$ (Yukawa=single number)

$$\to y_t \tilde{H} \boldsymbol{Q}\bar{\boldsymbol{u}} + y_t^* e^{i\theta_9} e^{-\frac{2\pi}{3\alpha_S(\Lambda_9)}} H \boldsymbol{Q}\bar{\boldsymbol{d}} + \frac{i3\theta_9}{32\pi^2} \int \left(G\tilde{G} + K\tilde{K}\right)$$ (Yukawa$\propto \mathbb{I}_3$, Flavor-diag)

$\bar{\theta} = -3\theta_9 + \arg\det|y_t|^2 \, e^{i\theta_9} = -3\theta_9 + 3\theta_9 = 0$ ✓

From now on, we set $\bar{\boldsymbol{\theta}}_{\boldsymbol{9}} = \mathbf{0}$ and take **real yuakwas.**

Outline

I. Generalized Global Symmetries

I-1. Higher-form symmetry
I-2. Non-invertible symmetry


II. Strong CP Problem-I: IR to UV

II-1. Non-invertible Peccei-Quinn symmetry
II-2. Massless quark solution

III. Strong CP Problem-II: UV to IR

III-1. $SU(9)$ Color-Flavor unification
III-2. Flavor structure and CKM CPV phase
III-3. Quality Problem

Solving Strong CP with Non-Invertible Symmetry



$SU(9)$

$\langle\Phi\rangle$

$SU(3)_C \times SU(3)_H/Z_3$

$\mathcal{L}_{UV} \supset y_u \tilde{H} Q \bar{u} + y_u^* e^{-\frac{2\pi}{\alpha_9}} H Q \bar{d} + \cdots$

with $\bar{\theta} = 0$

- Start with only $y_u \tilde{H} Q \bar{u}$ $(y_d = 0)$
- $Z_3^{\bar{d}}$ **NIS** from fractional CFU instantons
- $y_d H Q \bar{d}$ protected by **NIS**
- $\bar{\theta} =$ unphysical

$\langle\Sigma\rangle$

SM

Yukawa texture and CKM CPV (+possible $\bar{\theta}$)

with $\bar{\theta} = 0$ or $\bar{\theta} \ll 1$

Color-Flavor Unification

3. $SU(3)_C \times SU(3)_H/Z_3 \to SU(3)_C$: Flavor Structre and CKM CPV


(i) Two $SU(3)_H$ adjoint scalar $\Sigma_{1,2}$: $SU(3)_H \to \emptyset$

Complete breaking $\Leftrightarrow$ $[\Sigma_1, \Sigma_2] = \frac{[\Sigma^\dagger, \Sigma]}{2i} \neq 0$

(ii) Textures of y_u, y_d generated by structure of $\langle \Sigma_{1,2} \rangle$

Generate complex 3×3 y_u, y_d but in a way that $\bar{\theta} = 0$ is maintained.

$\Rightarrow$ **Our mechanism:** generate **Hermitian Yukawas**
(I) all CPV in scalar sector
(II) CPV transferred to SM fermions via bosonic mediation




$+$ diagram with $h.c.\left(\eta \leftrightarrow \eta^\dagger, \Sigma \leftrightarrow \Sigma^\dagger\right)$

Color-Flavor Unification

3. $SU(3)_C \times SU(3)_H/Z_3 \to SU(3)_C$: Flavor Structre and CKM CPV

(ii) Textures of y_u, y_d generated by structure of $\langle \Sigma_{1,2} \rangle$

Generate complex 3×3 y_u, y_d but in a way that $\bar{\theta} = 0$ is maintained.




$+$ diagram with $h.c.\left(\eta \leftrightarrow \eta^\dagger, \Sigma \leftrightarrow \Sigma^\dagger\right)$

Color-Flavor Unification

3. $SU(3)_C \times SU(3)_H/Z_3 \to SU(3)_C$: Flavor Structre and CKM CPV

(ii) Textures of y_u, y_d generated by structure of $\langle \Sigma_{1,2} \rangle$

Generate complex 3×3 y_u, y_d but in a way that $\bar{\theta} = 0$ is maintained.



$+$ diagram with $h.c.\left(\eta \leftrightarrow \eta^\dagger, \Sigma \leftrightarrow \Sigma^\dagger\right)$

$$(y_u)^i_j \sim y_t \left(1 + \frac{\alpha_9}{4\pi} \frac{\{\Sigma^\dagger, \Sigma\}}{2\Lambda_9^2} + \left(\frac{\alpha_9}{4\pi} \frac{\eta_1^\dagger (\Sigma^\dagger)^4 + \eta_2^\dagger (\Sigma^\dagger)^2 (\Sigma^\dagger)^2}{\Lambda_9^4} + h.c. \right) \right)^i_j$$


Hermitian **Real** **Complex**

Color-Flavor Unification

3. $SU(3)_C \times SU(3)_H/Z_3 \to SU(3)_C$: Flavor Structre and CKM CPV

(ii) Textures of y_u, y_d generated by structure of $\langle \Sigma_{1,2} \rangle$

Generate complex 3×3 y_u, y_d but in a way that $\bar{\theta} = 0$ is maintained.



$+$ diagram with $h.c.\left(\eta \leftrightarrow \eta^\dagger, \Sigma \leftrightarrow \Sigma^\dagger\right)$

$$(y_u)^i_j \sim y_t \left(1 + \frac{\alpha_9}{4\pi} \frac{\{\Sigma^\dagger, \Sigma\}}{2\Lambda_9^2} + \left(\frac{\alpha_9}{4\pi} \frac{\eta_1^\dagger (\Sigma^\dagger)^4 + \eta_2^\dagger (\Sigma^\dagger)^2 (\Sigma^\dagger)^2}{\Lambda_9^4} + h.c. \right) \right)^i_j$$

Hermitian **Real** **Complex**

$y_u, y_d \Rightarrow$ real eigenvalues $\Rightarrow$ $\bar{\theta} = \arg e^{-i\theta} \det(y_u y_d) = \arg \det(y_u y_d) = 0$ ✓

Color-Flavor Unification

3. $SU(3)_C \times SU(3)_H/Z_3 \to SU(3)_C$: Flavor Structre and CKM CPV

(iii) Generate $O(1)$ CKM CPV phase δ_{CKM} (without destabilizing $\boldsymbol{\overline{\theta} = 0}$)

Field-redefinition invariant definition of CKM CPV phase

$\tilde{J} = \mathrm{Im}\,\det\left[y_u^\dagger y_u, y_d^\dagger y_d\right] \propto \sin\delta_{CKM}$ "Jarlskog invariant"

Color-Flavor Unification

3. $SU(3)_C \times SU(3)_H/Z_3 \to SU(3)_C$: Flavor Structre and CKM CPV

(iii) Generate $O(1)$ CKM CPV phase δ_{CKM} (without destabilizing $\boldsymbol{\overline{\theta} = 0}$)

Field-redefinition invariant definition of CKM CPV phase

$$\tilde{J} = \mathrm{Im}\,\mathrm{det}\left[y_u^\dagger y_u, y_d^\dagger y_d\right] \propto \sin\delta_{CKM}$$

"Jarlskog invariant"

"misalignment" of y_u and y_d

Color-Flavor Unification

3. $SU(3)_C \times SU(3)_H/Z_3 \to SU(3)_C$: Flavor Structre and CKM CPV

(iii) Generate $O(1)$ CKM CPV phase δ_{CKM} (without destabilizing $\overline{\boldsymbol{\theta}} = \mathbf{0}$)

Field-redefinition invariant definition of CKM CPV phase


$$\tilde{J} = \mathrm{Im}\,\mathrm{det}\left[y_u^\dagger y_u, y_d^\dagger y_d\right] \propto \sin\delta_{CKM}$$ "Jarlskog invariant"

"misalignment" of y_u and y_d

So far, we have

$$y_u \sim y_t\left(1 + \{\Sigma^\dagger, \Sigma\} + (\eta\Sigma^4 + h.c.) + \cdots\right)$$

$$y_d \sim y_t^* e^{-\frac{2\pi}{\alpha_9}}\left(1 + \{\Sigma^\dagger, \Sigma\} + (\eta\Sigma^4 + h.c.) + \cdots\right)$$



Color-Flavor Unification

3. $SU(3)_C \times SU(3)_H/Z_3 \to SU(3)_C$: Flavor Structre and CKM CPV

(iii) Generate $O(1)$ CKM CPV phase δ_{CKM} (without destabilizing $\overline{\boldsymbol{\theta}} = \mathbf{0}$)

Field-redefinition invariant definition of CKM CPV phase

$$\tilde{J} = \mathrm{Im}\,\mathrm{det}\left[y_u^\dagger y_u, y_d^\dagger y_d\right] \propto \sin\delta_{CKM}$$


"Jarlskog invariant"

"misalignment" of y_u and y_d

So far, we have

$$y_u \sim y_t\left(1 + \{\Sigma^\dagger, \Sigma\} + (\eta\Sigma^4 + h.c.) + \cdots\right)$$

$$y_d \sim y_t^* e^{-\frac{2\pi}{\alpha_9}}\left(1 + \{\Sigma^\dagger, \Sigma\} + (\eta\Sigma^4 + h.c.) + \cdots\right)$$



Color-Flavor Unification

3. $SU(3)_C \times SU(3)_H/Z_3 \to SU(3)_C$: Flavor Structre and CKM CPV

(iii) Generate $O(1)$ CKM CPV phase δ_{CKM} (without destabilizing $\overline{\boldsymbol{\theta}} = \mathbf{0}$)

Field-redefinition invariant definition of CKM CPV phase

$$\tilde{J} = \mathrm{Im}\,\mathrm{det}\left[y_u^\dagger y_u, y_d^\dagger y_d\right] \propto \sin\delta_{CKM}$$

"Jarlskog invariant"

"misalignment" of y_u and y_d

So far, we have

$$y_u \sim y_t\left(1 + \{\Sigma^\dagger, \Sigma\} + (\eta\Sigma^4 + h.c.) + \cdots\right)$$

$$y_d \sim y_t^* e^{-\frac{2\pi}{\alpha_9}}\left(1 + \{\Sigma^\dagger, \Sigma\} + (\eta\Sigma^4 + h.c.) + \cdots\right)$$

$\boldsymbol{y_d \propto y_u}$ as a matrix

$$\Rightarrow\ \tilde{\boldsymbol{J}} = \mathbf{0}$$

Color-Flavor Unification

3. $SU(3)_C \times SU(3)_H/Z_3 \to SU(3)_C$: Flavor Structre and CKM CPV

(iii) Generate $O(1)$ CKM CPV phase δ_{CKM} (without destabilizing $\boldsymbol{\overline{\theta} = 0}$)

Field-redefinition invariant definition of CKM CPV phase

$$\tilde{J} = \mathrm{Im}\,\mathrm{det}\left[y_u^\dagger y_u, y_d^\dagger y_d\right] \propto \sin\delta_{CKM}$$

"Jarlskog invariant"

"misalignment" of y_u and y_d

So far, we have

$$y_u \sim y_t\left(1 + \{\Sigma^\dagger, \Sigma\} + (\eta\Sigma^4 + h.c.) + \cdots\right)$$

$$y_d \sim y_t^* e^{-\frac{2\pi}{\alpha_9}}\left(1 + \{\Sigma^\dagger, \Sigma\} + (\eta\Sigma^4 + h.c.) + \cdots\right)$$

$\boldsymbol{y_d \propto y_u}$ as a matrix

$\Rightarrow\ \boldsymbol{\tilde{J} = 0}$

We need extra ingredients to **misalign y_d vs y_u** : **'down-philic' interactions**

Color-Flavor Unification

3. $SU(3)_C \times SU(3)_H/Z_3 \to SU(3)_C$: Flavor Structre and CKM CPV

(iii) Generate $O(1)$ CKM CPV phase δ_{CKM} (without destabilizing $\overline{\boldsymbol{\theta}} = \mathbf{0}$)

	$SU(9)$	$U(1)_{Q-\bar{u}+\bar{d}}$
Φ	165 (3S)	0
$\Sigma_{1,2}$	80 (adj)	0
ρ	9	-1
χ	1	0

$$\mathcal{L}_{\rho\chi} \sim \lambda_d \bar{d}\rho\chi + \rho\left(c_1\Sigma^\dagger\Sigma + c_2\Sigma\Sigma^\dagger\right)\rho^\dagger$$

$$+ a_1\rho\Sigma\rho^\dagger + a_2\rho\Sigma\Sigma\rho^\dagger + h.c.$$

Color-Flavor Unification

3. $SU(3)_C \times SU(3)_H/Z_3 \to SU(3)_C$: Flavor Structre and CKM CPV

(iii) Generate $O(1)$ CKM CPV phase δ_{CKM} (without destabilizing $\overline{\boldsymbol{\theta}} = \mathbf{0}$)

	$SU(9)$	$U(1)_{Q-\bar{u}+\bar{d}}$
Φ	165 (3S)	0
$\Sigma_{1,2}$	80 (adj)	0
ρ	9	-1
χ	1	0

$$\mathcal{L}_{\rho\chi} \sim \lambda_d \bar{d}\rho\chi + \rho\left(c_1\Sigma^\dagger\Sigma + c_2\Sigma\Sigma^\dagger\right)\rho^\dagger$$
$$+ a_1\rho\Sigma\rho^\dagger + a_2\rho\Sigma\Sigma\rho^\dagger + h.c.$$

- Use χ rotation to set $\lambda_d \in \mathbb{R}$
- $c_{1,2} \in \mathbb{R}$
- $a_{1,2} \in \mathbb{C} \;\to\; a_1^2 a_2^\dagger, \eta_1^\dagger a_2^2$: new CPV source
- $a_{1,2} = 0$ if Z_4^Σ is imposed

  (again, our mechanism works regardless)


Color-Flavor Unification


3. $SU(3)_C \times SU(3)_H/Z_3 \to SU(3)_C$: Flavor Structre and CKM CPV

(iii) Generate $O(1)$ CKM CPV phase δ_{CKM} (without destabilizing $\overline{\boldsymbol{\theta}} = \mathbf{0}$)

$$\mathcal{L}_{\rho\chi} \sim \lambda_d \bar{d}\rho\chi + \rho\left(c_1\Sigma^\dagger\Sigma + c_2\Sigma\Sigma^\dagger\right)\rho^\dagger$$

$$+ a_1\rho\Sigma\rho^\dagger + a_2\rho\Sigma\Sigma\rho^\dagger + h.c.$$





Without "down-philic" interactions

$$y_u \sim y_t\left(1 + \{\Sigma^\dagger, \Sigma\} + (\eta\Sigma^4 + h.c.) + \cdots\right)$$

$$y_d \sim y_t^* e^{-\frac{2\pi}{\alpha_9}}\left(1 + \{\Sigma^\dagger, \Sigma\} + (\eta\Sigma^4 + h.c.) + \cdots\right)$$

$$\tilde{J} = 0$$


Color-Flavor Unification

3. $SU(3)_C \times SU(3)_H/Z_3 \to SU(3)_C$: Flavor Structre and CKM CPV

(iii) Generate $O(1)$ CKM CPV phase δ_{CKM} (without destabilizing $\bar{\theta} = 0$)

$$\mathcal{L}_{\rho\chi} \sim \lambda_d \bar{d}\rho\chi + \rho\left(c_1\Sigma^\dagger\Sigma + c_2\Sigma\Sigma^\dagger\right)\rho^\dagger$$


$$+ a_1\rho\Sigma\rho^\dagger + a_2\rho\Sigma\Sigma\rho^\dagger + h.c.$$



With "down-philic" interactions ($a_{1,2} = 0$)

$$y_u \sim y_t\left(1 + \{\Sigma^\dagger, \Sigma\} + (\eta\Sigma^4 + h.c.) + \cdots\right)$$

$$y_d \sim y_t^* e^{-\frac{2\pi}{\alpha_9}}\left(1 + \{\Sigma^\dagger, \Sigma\} + (\eta\Sigma^4 + h.c.) + \cdots + c\Sigma^\dagger\Sigma\right)$$



$$\tilde{J} \sim \mathrm{Im}\,\det\left(4r^2[\eta\Sigma^4 + \eta^\dagger\Sigma^{\dagger 4}, c\Sigma^\dagger\Sigma]\right), \qquad r \sim e^{-\frac{2\pi}{\alpha_9}}$$

$$\propto \mathrm{Im}\,\det\left(\eta\begin{pmatrix}[\Sigma,\Sigma^\dagger]\Sigma^4 + \Sigma[\Sigma,\Sigma^\dagger]\Sigma^3 + \Sigma^2[\Sigma,\Sigma^\dagger]\Sigma^2 \\ +\Sigma^3[\Sigma,\Sigma^\dagger]\Sigma\end{pmatrix} - h.c.\right)$$

Outline

I. Generalized Global Symmetries

I-1. Higher-form symmetry
I-2. Non-invertible symmetry

II. Strong CP Problem-I: IR to UV

II-1. Non-invertible Peccei-Quinn symmetry
II-2. Massless quark solution

III. Strong CP Problem-II: UV to IR


III-1. $SU(9)$ Color-Flavor unification
III-2. Flavor structure and CKM CPV phase
III-3. Quality Problem

No/Less Quality Problem

1. Our solution requires "high-quality" $U(1)_{PQ}$ in the UV

No/Less Quality Problem

1. Our solution requires "high-quality" $U(1)_{PQ}$ in the UV



$= 9\ (\square)\ \text{of}\ SU(9)$

	$SU(9)$	$U(1)_{Q-\bar{u}}$	$U(1)_{\bar{d}}$
$\boldsymbol{Q} = (\boldsymbol{u}, \boldsymbol{d})^{\boldsymbol{t}}$	9	$+1$	0
$\bar{\boldsymbol{u}}$	$\bar{9}$	-1	0
$\bar{\boldsymbol{d}}$	$\bar{9}$	0	$+1$
H	1	0	0

$$\mathcal{L}_0 = y_t \tilde{H} \boldsymbol{Q} \bar{\boldsymbol{u}} + h.c. + \frac{i\theta_9}{32\pi^2} \int F\tilde{F}$$

$U(1)_{Q-\bar{u}}[SU(9)]^2 = U(1)_{\bar{d}}[SU(9)]^2 = 1$

⇒ [Anomaly Free] $U(1)_{B=Q-\bar{u}-\bar{d}}$

[Anomalous] $U(1)_{Q-\bar{u}+\bar{d}}$ or $U(1)_{\bar{d}}$


No/Less Quality Problem

1. Our solution requires "high-quality" $U(1)_{PQ}$ in the UV

$SU(3)_H$ →, $SU(3)_C$ ↓

$\bar{u}$	$\bar{c}$	$\bar{t}$
$\bar{u}$	$\bar{c}$	$\bar{t}$
$\bar{u}$	$\bar{c}$	$\bar{t}$

$= 9\ (\square)$ of $SU(9)$



	$SU(9)$	$U(1)_{Q-\bar{u}}$	$U(1)_{\bar{d}}$
$\boldsymbol{Q} = (\boldsymbol{u}, \boldsymbol{d})^{\boldsymbol{t}}$	9	$+1$	0
$\bar{\boldsymbol{u}}$	$\bar{9}$	-1	0
$\bar{\boldsymbol{d}}$	$\bar{9}$	0	$+1$
H	1	0	0

$$\mathcal{L}_0 = y_t \tilde{H} \boldsymbol{Q}\bar{\boldsymbol{u}} + h.c. + \frac{i\theta_9}{32\pi^2}\int F\tilde{F}$$

$$+ y_t^* e^{i\theta_9} e^{-\frac{2\pi}{\alpha_9} H \boldsymbol{Q}\bar{\boldsymbol{d}}}$$

$U(1)_{Q-\bar{u}}[SU(9)]^2 = U(1)_{\bar{d}}[SU(9)]^2 = 1$

⇒ [Anomaly Free] $U(1)_{B=Q-\bar{u}-\bar{d}}$

[Anomalous] $U(1)_{Q-\bar{u}+\bar{d}}$ or $U(1)_{\bar{d}}$

No/Less Quality Problem


1. Our solution requires "high-quality" $U(1)_{PQ}$ in the UV

2. Estimation of Needed Quality from Quantum Gravity Effects

No/Less Quality Problem

1. Our solution requires "high-quality" $U(1)_{PQ}$ in the UV

2. Estimation of Needed Quality from Quantum Gravity Effects




$$y_d = 0$$


No/Less Quality Problem

1. Our solution requires "high-quality" $U(1)_{PQ}$ in the UV

2. Estimation of Needed Quality from Quantum Gravity Effects



$y_d = 0$




$y_t^* e^{i\theta_9} e^{-\frac{2\pi}{\alpha_9} H \boldsymbol{Q}\bar{\boldsymbol{d}}}$


No/Less Quality Problem

1. Our solution requires "high-quality" $U(1)_{PQ}$ in the UV


2. Estimation of Needed Quality from Quantum Gravity Effects



$y_d = 0$



$y_t^* e^{i\theta_9} e^{-\frac{2\pi}{\alpha_9} H \boldsymbol{Q}\overline{\boldsymbol{d}}}$




QG breaking of $U(1)_{PQ}$ with $O(1)$ phase


No/Less Quality Problem

1. Our solution requires "high-quality" $U(1)_{PQ}$ in the UV


2. Estimation of Needed Quality from Quantum Gravity Effects




$y_d = 0$



$y_t^* e^{i\theta_9} e^{-\frac{2\pi}{\alpha_9} H \boldsymbol{Q}\bar{\boldsymbol{d}}}$



QG breaking of $U(1)_{PQ}$ with $O(1)$ phase




$\bar{\theta} \ll 1$ if

$|\text{QG effect}| \ll |SU(9)\text{inst}|$


No/Less Quality Problem

1. Our solution requires "high-quality" $U(1)_{PQ}$ in the UV


2. Estimation of Needed Quality from Quantum Gravity Effects




$y_d = 0$



$y_t^* e^{i\theta_9} e^{-\frac{2\pi}{\alpha_9} H \boldsymbol{Q}\bar{\boldsymbol{d}}}$



QG breaking of $U(1)_{PQ}$ with $O(1)$ phase



$\bar{\theta} \ll 1$ if


$|\text{QG effect}| \ll |SU(9)\text{inst}|$

$$\frac{|\Phi|^2 HQ\bar{d}}{M_{\text{Pl}}^2} \rightarrow \langle\Phi\rangle \sim \Lambda_9 < 10^{13}\ \text{GeV}$$

$$\frac{HQ\Sigma\bar{d}}{M_{\text{Pl}}} \rightarrow \langle\Sigma\rangle \sim \Lambda_3 < 10^{8}\ \text{GeV}$$


Conclusion


Solving Strong CP with Non-Invertible Symmetry



- Start with only $y_u \tilde{H} Q \bar{u} \quad (y_d = 0)$
- $Z_3^{\bar{d}}$ **NIS** from fractional CFU instantons ($\boldsymbol{N_C = N_g}$)
- $y_d H Q \bar{d}$ protected by **NIS**
- $\bar{\theta} =$ unphysical

Solving Strong CP with Non-Invertible Symmetry







Small (G/H) Instanton

$$\mathcal{L}_{UV} \supset y_u \tilde{H} Q \bar{u} + y_u^* e^{-\frac{2\pi}{\alpha_9}} H Q \bar{d} + \cdots$$

- Start with only $y_u \tilde{H} Q \bar{u} \quad (y_d = 0)$
- $Z_3^{\bar{d}}$ **NIS** from fractional CFU instantons ($\boldsymbol{N_C = N_g}$)
- $y_d H Q \bar{d}$ protected by **NIS**
- $\bar{\theta} =$ unphysical

Solving Strong CP with Non-Invertible Symmetry





Small (G/H) Instanton

$$\mathcal{L}_{UV} \supset y_u \tilde{H} Q \bar{u} + y_u^* e^{-\frac{2\pi}{\alpha_9}} H Q \bar{d} + \cdots$$

- Start with only $y_u \tilde{H} Q \bar{u} \quad (y_d = 0)$
- $Z_3^{\bar{d}}$ **NIS** from fractional CFU instantons ($\boldsymbol{N_C = N_g}$)
- $y_d H Q \bar{d}$ protected by **NIS**
- $\bar{\theta} =$ unphysical

Yukawa texture and CKM CPV (+possible $\bar{\theta}$)
(Hermitian Yukawa + down-philic interactions)

THANK YOU

FOR

YOUR ATTENTION!


Non-Invertible Symmetry

1. From $U(1)$ Instanton

Half-spacing gauging construction

$$\mathcal{T}(\theta)$$


$$S = \frac{1}{4e^2}\int F_2 \wedge * F_2 + \frac{i\theta}{8\pi^2}\int F_2 \wedge F_2$$



Non-Invertible Symmetry

1. From $U(1)$ Instanton

Half-spacing gauging construction



$$S = \frac{1}{4e^2}\int F_2 \wedge * F_2 + \frac{i\theta}{8\pi^2}\int F_2 \wedge F_2$$


[Gauge $Z_N^{(1)}(m) \subset U(1)^{(1)}(m)$]

$$S = \frac{1}{4e^2}\int F_2 \wedge * F_2 + \frac{i\theta}{8\pi^2}\int F_2 \wedge F_2 + \frac{i}{2\pi}\int B_2 \wedge F_2 + \frac{iN}{2\pi}\int B_2 \wedge dC_1 + \frac{ikN}{4\pi}\int B_2 \wedge B_2$$

Non-Invertible Symmetry

1. From $U(1)$ Instanton

Half-spacing gauging construction



$\mathcal{T}(\theta)$

$$S = \frac{1}{4e^2}\int F_2 \wedge * F_2 + \frac{i\theta}{8\pi^2}\int F_2 \wedge F_2$$

[Gauge $Z_N^{(1)}(m) \subset U(1)^{(1)}(m)$]


$$\mathcal{T}\left(\theta - \frac{2\pi p}{N}\right) \quad (kp=1 \bmod N)$$

$$S = \frac{1}{4e^2}\int F_2 \wedge * F_2 + \frac{i\theta}{8\pi^2}\int F_2 \wedge F_2 + \frac{i}{2\pi}\int B_2 \wedge F_2 + \frac{iN}{2\pi}\int B_2 \wedge dC_1 + \frac{ikN}{4\pi}\int B_2 \wedge B_2$$

Non-Invertible Symmetry

1. From $U(1)$ Instanton

Half-spacing gauging construction



$\mathcal{T}(\theta)$

$$S = \frac{1}{4e^2}\int F_2 \wedge * F_2 + \frac{i\theta}{8\pi^2}\int F_2 \wedge F_2$$

[Gauge $Z_N^{(1)}(m) \subset U(1)^{(1)}(m)$]

$$\mathcal{T}\left(\theta - \frac{2\pi p}{N}\right) \quad (kp=1 \bmod N)$$

$$S = \frac{1}{4e^2}\int F_2 \wedge * F_2 + \frac{i\theta}{8\pi^2}\int F_2 \wedge F_2 + \frac{i}{2\pi}\int B_2 \wedge F_2 + \frac{iN}{2\pi}\int B_2 \wedge dC_1 + \frac{ikN}{4\pi}\int B_2 \wedge B_2$$

[Anomalous Field Redefinition]


$\mathrm{U}(1)_\mathrm{A}$: $\Psi \to e^{2\pi i/z}\,\Psi$

$$S \to S + \frac{2\pi i}{z}\int d * j_1 + \frac{2\pi i K}{z}\int \frac{F\wedge F}{8\pi^2} \quad \leftrightarrow \quad \left[\mathcal{T}(\theta) \to \mathcal{T}\left(\theta + \frac{2\pi K}{z}\right)\right]$$

Non-Invertible Symmetry

1. From $U(1)$ Instanton

Half-spacing gauging construction



$$S = \frac{1}{4e^2}\int F_2 \wedge * F_2 + \frac{i\theta}{8\pi^2}\int F_2 \wedge F_2$$

[Gauge $Z_N^{(1)}(m) \subset U(1)^{(1)}(m)$]


[Anomalous Field Redefinition]

$$\frac{p}{N} = \frac{K}{z}$$

Non-Invertible Symmetry

1. From $U(1)$ Instanton

Half-spacing gauging construction



$\mathcal{T}(\theta)$

$$S = \frac{1}{4e^2}\int F_2 \wedge * F_2 + \frac{i\theta}{8\pi^2}\int F_2 \wedge F_2$$

[Gauge $Z_N^{(1)}(m) \subset U(1)^{(1)}(m)$]

$$\mathcal{T}\left(\theta - \frac{2\pi p}{N}\right) \quad (kp=1 \bmod N)$$

$$S = \frac{1}{4e^2}\int F_2 \wedge * F_2 + \frac{i\theta}{8\pi^2}\int F_2 \wedge F_2 + \frac{i}{2\pi}\int B_2 \wedge F_2 + \frac{iN}{2\pi}\int B_2 \wedge dC_1 + \frac{ikN}{4\pi}\int B_2 \wedge B_2$$

[Anomalous Field Redefinition]

$\mathrm{U}(1)_\mathrm{A}$: $\Psi \to e^{2\pi i/z}\,\Psi$

$$S \to S + \frac{2\pi i}{z}\int d * j_1 + \frac{2\pi i K}{z}\int \frac{F\wedge F}{8\pi^2} \quad \leftrightarrow \quad \left[\mathcal{T}(\theta) \to \mathcal{T}\left(\theta + \frac{2\pi K}{z}\right)\right]$$


Non-Invertible Symmetry

1. From $U(1)$ Instanton

Half-spacing gauging construction

$$\mathcal{T}(\theta)$$

$$S = \frac{1}{4e^2}\int F_2 \wedge * F_2 + \frac{i\theta}{8\pi^2}\int F_2 \wedge F_2$$



[Anomalous Field Redefinition]

$\mathrm{U}(1)_\mathrm{A}$: $\Psi \to e^{2\pi i/z}\,\Psi$

$$S \to S + \frac{2\pi i}{z}\int d * j_1 + \frac{2\pi i K}{z}\int \frac{F\wedge F}{8\pi^2} \quad \leftrightarrow \quad \left[\mathcal{T}(\theta) \to \mathcal{T}\left(\theta + \frac{2\pi K}{z}\right)\right]$$

Non-Invertible Symmetry

1. From $U(1)$ Instanton


Half-spacing gauging construction

$\mathcal{T}(\theta)$

$$S = \frac{1}{4e^2}\int F_2 \wedge * F_2 + \frac{i\theta}{8\pi^2}\int F_2 \wedge F_2$$

[Gauge $Z_N^{(1)}(m) \subset U(1)^{(1)}(m)$]

$$S = \frac{1}{4e^2}\int F_2 \wedge * F_2 + \frac{i\theta}{8\pi^2}\int F_2 \wedge F_2 + \frac{i}{2\pi}\int B_2 \wedge F_2 + \frac{iN}{2\pi}\int B_2 \wedge dC_1 + \frac{ikN}{4\pi}\int B_2 \wedge B_2$$



[Anomalous Field Redefinition]

$\mathrm{U(1)_A}$: $\Psi \to e^{2\pi i/z}\,\Psi$

$$S \to S + \frac{2\pi i}{z}\int d * j_1 + \frac{2\pi i K}{z}\int \frac{F\wedge F}{8\pi^2} \quad \leftrightarrow \quad \left[\mathcal{T}(\theta) \to \mathcal{T}\left(\theta + \frac{2\pi K}{z}\right)\right]$$

Non-Invertible Symmetry

1. From $U(1)$ Instanton


Half-spacing gauging construction

$\mathcal{T}(\theta)$

$$S = \frac{1}{4e^2}\int F_2 \wedge * F_2 + \frac{i\theta}{8\pi^2}\int F_2 \wedge F_2$$

[Gauge $Z_N^{(1)}(m) \subset U(1)^{(1)}(m)$]

$$S = \frac{1}{4e^2}\int F_2 \wedge * F_2 + \frac{i\theta}{8\pi^2}\int F_2 \wedge F_2 + \frac{i}{2\pi}\int B_2 \wedge F_2 + \frac{iN}{2\pi}\int B_2 \wedge dC_1 + \frac{ikN}{4\pi}\int B_2 \wedge B_2$$



[Anomalous Field Redefinition]


$\mathrm{U(1)_A}$: $\Psi \to e^{2\pi i/z}\,\Psi$

$$S \to S + \frac{2\pi i}{z}\int d * j_1 + \frac{2\pi i K}{z}\int \frac{F\wedge F}{8\pi^2} \quad \leftrightarrow \quad \left[\mathcal{T}(\theta) \to \mathcal{T}\left(\theta + \frac{2\pi K}{z}\right)\right]$$

Non-Invertible Symmetry

1. From $U(1)$ Instanton

Half-spacing gauging construction



[Anomalous Symmetry]

$\mathrm{U}(1)_{\mathrm{A}}$: $\Psi \to e^{2\pi i/z}\,\Psi$

$S \to S + \frac{2\pi i K}{z}\int \frac{F\wedge F}{8\pi^2}$


Non-Invertible Symmetry

1. From $U(1)$ Instanton

Half-spacing gauging construction

$$\mathcal{T}(\theta)$$

$$S = \frac{1}{4e^2}\int F_2 \wedge * F_2 + \frac{i\theta}{8\pi^2}\int F_2 \wedge F_2$$



[Anomalous Symmetry] × [$\mathcal{A}^{N,p}(F_2/N)$]


$\mathrm{U}(1)_\mathrm{A}$: $\Psi \to e^{2\pi i/z}\,\Psi$

$$S \to S + \frac{2\pi i K}{z}\int \frac{F\wedge F}{8\pi^2} - \frac{2\pi i p}{N}\int \frac{F\wedge F}{8\pi^2} \to S$$

Non-Invertible Symmetry

1. From $U(1)$ Instanton

Half-spacing gauging construction




[Anomalous Symmetry] × $[\mathcal{A}^{N,p}(F_2/N)]$

$\mathrm{U}(1)_\mathrm{A}$: $\Psi \to e^{2\pi i/z}\,\Psi$

$S \to S + \frac{2\pi i K}{z}\int \frac{F\wedge F}{8\pi^2} - \frac{2\pi i p}{N}\int \frac{F\wedge F}{8\pi^2} \to S$

Strong-CP Problem

1. Strong CP Problem



Expectation based on general rules of effective field theory

$$S_{QCD} \supset \frac{i\bar{\theta}}{8\pi^2}\int Tr(G \wedge G)$$

Neutron Electric Dipole Moment

$$d_n \sim 3 \times 10^{16}\,\bar{\theta} \;\rightarrow\; \bar{\theta} < 10^{-10}$$

$\tilde{J} = \mathbf{Im}\,\mathbf{det}[y_u^\dagger y_u, y_d^\dagger y_d] \propto \sin\delta_{CKM}$ "Jarlskog invariant"

vs

$\bar{\theta} \equiv \arg e^{-i\theta}\,\det(y_u y_d)$

Color-Flavor Unification

3. $SU(3)_C \times SU(3)_H/Z_3 \to SU(3)_C$: Flavor Structre and CKM CPV

	$SU(9)$	$U(1)_{Q-\bar{u}+\bar{d}}$
Φ	165 (3S)	0
$\Sigma_{1,2}$	80 (adj)	0
ρ	9	-1
χ	1	0

(i) Two $SU(3)_H$ adjoint scalar $\Sigma_{1,2}$: $SU(3)_H \to \emptyset$

(ii) Textures of y_u, y_d generated by structure of $\langle\Sigma_{1,2}\rangle$

(iii) Required CKM CPV phase from $V(\Sigma)$

Color-Flavor Unification

3. $SU(3)_C \times SU(3)_H/Z_3 \to SU(3)_C$: Flavor Structre and CKM CPV

	$SU(9)$	$U(1)_{Q-\bar{u}+\bar{d}}$
Φ	165 (3S)	0
$\Sigma_{1,2}$	80 (adj)	0
ρ	9	-1
χ	1	0

(i) Two $SU(3)_H$ adjoint scalar $\Sigma_{1,2}$: $SU(3)_H \to \emptyset$

(ii) Textures of y_u, y_d generated by structure of $\langle\Sigma_{1,2}\rangle$

(iii) Required CKM CPV phase from $V(\Sigma)$

Combine $\Sigma = \Sigma_1 + i\Sigma_2$ $(U(1)_\Sigma)$

Consider a simple case with Z_4 invariant potential $V(\Sigma)$
(our mechanism works regardless of this simplifying assumption)

$$V(\Sigma) = \eta_1 \mathrm{Tr}(\Sigma^4) + \eta_2 \left(\mathrm{Tr}(\Sigma^2)\right)^2 + h.c. + \xi \mathrm{Tr}\left(\Sigma^\dagger \Sigma\right)^2 + \cdots \text{(terms with real coeffs)}$$

Color-Flavor Unification

3. $SU(3)_C \times SU(3)_H/Z_3 \to SU(3)_C$: Flavor Structre and CKM CPV

	$SU(9)$	$U(1)_{Q-\bar{u}+\bar{d}}$
Φ	165 (3S)	0
$\Sigma_{1,2}$	80 (adj)	0
ρ	9	-1
χ	1	0

(i) Two $SU(3)_H$ adjoint scalar $\Sigma_{1,2}$: $SU(3)_H \to \emptyset$

(ii) Textures of y_u, y_d generated by structure of $\langle\Sigma_{1,2}\rangle$

(iii) Required CKM CPV phase from $V(\Sigma)$

Combine $\Sigma = \Sigma_1 + i\Sigma_2$ $(U(1)_\Sigma)$

Consider a simple case with Z_4 invariant potential $V(\Sigma)$
(our mechanism works regardless of this simplifying assumption)

$$V(\Sigma) = \eta_1 \mathrm{Tr}(\Sigma^4) + \eta_2 \left(\mathrm{Tr}(\Sigma^2)\right)^2 + h.c. + \xi \mathrm{Tr}\left(\Sigma^\dagger \Sigma\right)^2 + \cdots \text{(terms with real coeffs)}$$

Field redefinition invariant CPV : $\boldsymbol{\eta_1^\dagger \eta_2}$

$$\Sigma \to e^{-i\varphi_1/4}\Sigma: \quad |\eta_1| e^{i\varphi_1} \mathrm{Tr}(\Sigma^4) + |\eta_2| e^{i\varphi_2} \left(\mathrm{Tr}(\Sigma^2)\right)^2 \to |\eta_1| \mathrm{Tr}(\Sigma^4) + |\eta_2| e^{i(\varphi_2 - \varphi_1)} \left(\mathrm{Tr}(\Sigma^2)\right)^2$$

Color-Flavor Unification

3. $SU(3)_C \times SU(3)_H/Z_3 \to SU(3)_C$ **: Flavor Structre and CKM CPV**

(i) Two $SU(3)_H$ adjoint scalar $\Sigma_{1,2}$: $SU(3)_H \to \emptyset$

Complete breaking $\Leftrightarrow$ $[\Sigma_1, \Sigma_2] = \frac{[\Sigma^\dagger, \Sigma]}{2i} \neq 0$

(ii) Textures of y_u, y_d generated by structure of $\langle \Sigma_{1,2} \rangle$

Color-Flavor Unification

3. $SU(3)_C \times SU(3)_H/Z_3 \to SU(3)_C$: Flavor Structre and CKM CPV

(i) Two $SU(3)_H$ adjoint scalar $\Sigma_{1,2}$: $SU(3)_H \to \emptyset$

Complete breaking $\Leftrightarrow$ $[\Sigma_1, \Sigma_2] = \frac{[\Sigma^\dagger, \Sigma]}{2i} \neq 0$

(ii) Textures of y_u, y_d generated by structure of $\langle \Sigma_{1,2} \rangle$

Generate complex 3×3 y_u, y_d but in a way that $\bar{\theta} = 0$ is maintained.

$\Rightarrow$ **Our mechanism:** generate **Hermitian Yukawas**

(I) all CPV in scalar sector

(II) CPV transferred to SM fermions via bosonic mediation